







ACUTE CARE – PRIMARY SURVEY – “LOOK, HEAR & FEEL”

	Problem	Look /Listen/Feel	Intervene
Airway 	Direct trauma - Disruption - Oedema (later sign due to disruption)  Obstruction - FB - Food vomitus - Bloody Vomitus - Soft Tissue oedema (haematoma)  Other reasons for airway  - Decreased LOC - SEVERE lung pathology - CO poisoning - Surgery (RSI = Rapid sequence induction) to prevent risk of aspiration pneumonia + increase FRC	Appearance (the sweaty tachypnoic) Colour (look grey, pale) Posture (LOC) C-spine immobilise Airway obstruction signs - Stridor – swelling in larynx - hoarse voice Signs of imminent airway collapse (e.g. severe trauma, burn) - Protruding tongue - Drooling - Trismus - Hypoxia (late sign) Deteriorating consciousness Choking 5 back blows 5 chest blows	Position Upright (best) at least 30° to optimise breathing L) lateral position if unconscious Prepare + SUCTION → CLEAR AIRWAY - Gloved finger + suction Adjunct airway – centre of mouth to angle of jaw Manoeuvres – Chin lift/jaw thrust or neutral in infants - Oropharyngeal (guedel)/Nasopharyngeal airway Intubation → laryngoscope (CI = if NOT fasted) - Straighten the airway - LMA or ETT (8.0mm standard for adult males) Surgical Airway (Can't intubate, ventilate, LMA) - Cricothyroidotomy OR Tracheostomy Maintain in-line immobilisation (C-spine) - If LOC / head injuries - Manually, sandbags, cervical collar
Breathing Tachypnoea = 1st sign of sepsis and metabolic acidosis 	Massive flail ribs (part of ribs separated from rest of chest wall → ↓↓ LOC/poor resp effort) Simple pneumothorax Massive haemothorax Tension pneumothorax → impaired VR (compresses IVC = distended neck veins) Open Pneumothorax High SCI OR TAMPONADE 	Colour – Cyanosis+ Diaphoresis Chest expansion → asymmetry and reduced Accessory muscle + posture ↑WOB + ↑RR + ↓Sats (100% sats = CO, methamoglobin) Tracheal dev. + abdo breath + distended neck veins Speaking in complete sentences Noisy breathing (stridor, wheeze) Percussion (dullness, hyper-resonance, surgical emphysema) Auscultate (BS, Creps, wheeze) 	Recheck vitals + trends + meds Max FIO2 → mech. Ventilation - If in doubt = 4L/min Hudson mask (titrate after) Needle aspiration → 2nd IC in MCL Tube thoracostomy: 3-bottle chest drain → 5th IC in MAL → hug lower 6th rib (avoid neurovasbundle in costal groove superiorly) Open pneumothorax → Cover open wound (3 side occlusive) dressing to let air out but not in 
Circulation “stop bleeding” 	Bleeding (Big 5) - External (obs) - Chest (CXR) - Abdomen (FAST, DPL) - Pelvis (PXR) - Femurs (Clin. Exam) *Combination **DPL = diagnostic peritoneal lavage  Heart - Tension pneumothorax - Pericardial tamponade - Contusion - Infarction 	Colour (pale, sweaty) Warm (distributive), cold (other) HR, BP, CRT, JVP Urine output Fluid overload signs (raised JVP or distended neck veins, peripheral oedema) Dehydration signs (tachycardia, hypotTN, ↑RR, dry MM, reduced skin turgor, altered LOC) Peripheral Pulse – quality, regularity - Palpate apex beat (best for HF) Auscultate (HSDNM) Microcirculation - Skin – warmth? - Renal – oliguria? - Brain – hypoxia?LOC?	1st line = 2x IV or IO access –14-16g cannula Bloods = cultures FBC, Coags (DIC), X-match, ABG IV ABx empirical IVF = crystalloids (0.9% NS 500mL bolus), IVF colloids (pRBC – O neg) Pelvic stabiliser Needle thoracostomy / pericardiocentesis ED thoracostomy (REBOA) – catheter via femoral vein to stop inferior blood flow via balloon inflation Transfer to ICU or monitor on ward - Inotropes / vasopressors Long-term: Monitor Fluid balance INPUT (IV)= PO/IVF, NGT OUTPUT = IDC, NGT, drains, tubes Monitor EUC 
Disability 	2° brain injury (due to inadequate oxygenation to brain tissue) → can amplify damage caused by 1° brain injury INTRA= haematoma, oedema, fitting Extra – hypoxia, hypotTN, ↓/↑ CO2, BSL 	AVPU → GCS ≤ 8 (intubate) Cognition/ behaviour = orientation to place, person and time. Eyes = PEARL, nystagmus, ptosis Speech = slurred Motor (power/ tone) / sensation = Facial asymmetry, seizures, abnormal or absent limb movement GLUCOSE CHECK med chart (insulin, OHA) Hypo Sx (confusion, low BSL, polydipsia, disorientated) Diaphoresis (Sweaty, cold, clammy)	Position - C-spine protection - Controlled ventilation – prone position - Craniotomy (e.g. EDH, SDH) + Neurosurg - Closed reduction OR realign fractures - Debride ischaemic or contaminated wound Medications - For sedation - Analgesia - Anti-emetics
Exposure 	Hypothermia Rashes – location, distribution,, blanching? Thrombin (COAGS) – hidden bleeds - Check wounds, drains, IDCs 	Take OFF everything Prepare for 2° survey Monitor Temp 	Remove all clothes - Rashes - Skin appearance - Abdo exam Warm fluids, blankets. Heat mattress = bair hugger -3M Increase room temp.
Fluid status Glucose Hardware / lines Investigations	Urine Output and Glucose Beside = ECG (arrythmia) Urine dipstick (UTI, DKA), CXR (pneumothorax), pelvis X-ray, (PELVIC bleed) Bloods = ABG (check lactates, base excess – cellular function = ischaemia), IDC (UO), NGT (decompress bowel in SBO/LBO) FAST/eFAST (fcosu assessment w/ sonography in trauma → air/blood in pleural cavity + blood in abdo cavity + heart/liver) 2nd Survey → ABx, tetanus prophylaxis		

Advanced Life Support for Adults



During CPR
Airway adjuncts (LMA / ETT)
Oxygen
Waveform capnography
IV / IO access
Plan actions before interrupting compressions (e.g. charge manual defibrillator)

Drugs
Shockable
* Adrenaline 1 mg after 2nd shock (then every 2nd loop)
* Amiodarone 300mg after 3 shocks
Non Shockable
* Adrenaline 1 mg immediately (then every 2nd loop)

Consider and Correct
Hypoxia
Hypovolaemia
Hyper / hypokalaemia / metabolic disorders
Hypothermia / hyperthermia
Tension pneumothorax
Tamponade
Toxins
Thrombosis (pulmonary / coronary)

Post Resuscitation Care
Re-evaluate ABCDE
12 lead ECG
Treat precipitating causes
Aim for: SpO₂ 94-98%, normocapnia and normoglycaemia
Targeted temperature management

New points:

- C → B → A (NOT abc)
- Cycles are 2 minutely (AVOID interruptions - COACHED)
- **CPR is the MOST IMPORTANT & PROVEN Rx**
 - 30:2 (adults), 15:2 (child) – rate is critical for sufficient preload and cardiac output
 - **Fun Fact:** cough before going into arrest - to increase afterload
- **Early defibrillation is key** – 200J (adult), 4J/kg (child)
- **AVOID XS ventilation**
- **1mg Adrenaline** (after 2nd shock) at 4mins
 - **Nb:** Lower dosage of adrenaline used in anesthetics (25mcg, 50mcg – since potent vasoconstrictor)

Post-resus care (stabilise and transfer care)

- Team-based approach
 - 2nd survey = don't forget spine injuries
 - AMPLE history + collateral (bystander, ambo)
 - **Bloods** = FBC, EUC, LFT, Blood/Urine cultures,
 - **Bedside** = ECG, CXR, Urine dipstick, BSL, ABG, FAST
 - **Imaging** – CT brain/chest/abdo
 - **Identify and treat causes**
- Keep stable**
- **Analgesia** (intranasal fentanyl (child/adult) or morphine (adults))
 - **Sedation** (e.g. midazolam, propofol)
 - Maintain O₂ at 94-98%
 - Central Temp, glucose, and CO₂ (capnography)

What's the worst thing? = 1st dx

AAA, PE, ACS, aortic dissection, GI perforation, tension pneumothorax
➢ Always come up with 3 DDX




AIRWAY (LEMON)

L – Look externally for characteristics known to cause difficult laryngoscopy (please circle all that apply)

- Face**
- ☐ Small jaw
 - ☐ Facial hair
 - ☐ Difficult Bag/Mask Ventilation (2 person, use of airway, inability to maintain seal)
 - ☐ Edema
 - ☐ Prominent Teeth
 - ☐ Loose Teeth
 - ☐ Disfiguring of the Jaw
- Thorax / Abdomen**
- ☐ Pregnancy
 - ☐ Bowel Obstruction
 - ☐ Massive ascities
 - ☐ Morbid obesity


E – Evaluate the 3-3 Rule:

- Mouth opening – 3 finger breadths ☐ yes ☐ no
Thyro-Mental distance – 3 finger breadths ☐ yes ☐ no



M – Mallampati Score

Mallampati Class: _____



O – Obstruction (Is there any condition that can cause obstruction of the airway which would make laryngoscopy and ventilation difficult?)

- ☐ Tumors
- ☐ Stridor
- ☐ Congenital Defects (Down's, Goiter, Pierre-Robin Syndrome)
- ☐ Other obvious deformity

N – Neck mobility

- Can the patient move their jaw forward? ☐ yes ☐ no
Can the patient fully bend / extend the head and neck? ☐ yes ☐ no
Is the patient in a cspine collar? ☐ yes ☐ no

Assessment

- History:** AMPLE
- Exam:** LEMON, LOC, haemodynamics
(**LEMON = should we intubate?**)
- Risk factors** for difficult airway
 - Trauma, High BMI
 - FB, Congenital tumour
 - Burns

Management

- Least invasive
- Basic manoeuvres** (chin lift with head tilt, or jaw thrust – better for c-spine issue)
AIM to improve axis – straighten airway
- Add adjuncts** (Guedel, NPA, LMA)
- BVM – **2 person technique**
- Advanced techniques (I+V)

Vortex

- After best attempt at any airway method (e.g. LMA, BVM or Intubation)
- AIM = stay out of vortex
- Go for cricothyroidotomy (surgical airway)

NB: good sats may NOT mean a good airway



	Airway Adjuncts			Intubation	Surgical airway
	Guedel (OPA)	NPA	LMA	I+V	Tracheotomy
Ind	ALS	ALS	Anaesthesia Resus	1) Create airway (blockage) 2) Maintain airway (prevent) 3) Protect airway (LOC) 4) Ventilate (if lung pathology)	➢ Can't intubate, ventilate, LMA ➢ CI = futile airway, if able to secure airway less invasively
Mx	Angle of jaw to incisors	➢ Nostril to ear lobe ➢ Lubricate ➢ Can cause blood loss (esp. patient on aspirin)	Choose correct size (8mm -males, 7mm - females) ➢ Lubricate		➢ Cricothyroidotomy ➢ Tracheotomy

PATIENT SAFETY AND QUALITY SCENARIOS

ALS in ED

SCENARIO

1. Arrest called at triage - Alert alarm at triage, everyone goes running
 - 60 year old man, walked into ED with wife, arrested in triage area
2. Arrest on arrival into resus - 68 year old man BIBA with severe SOB
 - Arrested moments after transfer onto resus bed
3. Bat call – 50 year old female, arrested at a swimming pool
 - CPR in progress with ambulance, 2 x Defib delivered so far
4. Code blue call to radiology – 42 year old female found arrested by staff in waiting bay
 - Inpatient, 20 minutes post CTPA, was in a waiting area for transfer back to ward

*Notes on ED – loud, lots of staff if day time, not enough at night, may need crowd control, team work
Often little is known about the patient*

ALS in Anaesthetics

SCENARIO

1. 31 year old male for laparoscopic gastric sleeve. 110kg otherwise well
 - Induction for anaesthetic – midazolam, fentanyl, propofol, rocuronium
 - Severe bronchospasm, difficult to ventilate, BP 50/30
 - Metaraminol, intubated – no CO2, reintubated with CMAC still no CO2. 100% oxygen
 - Adrenaline boluses then infusion, CPR, hand bagged, non shockable
 - VF, shock, ECMO 45 min post arrest, output restored
 - TF to major ECMO centre

Notes on Anaesthetics – Relatively rare, usually lots of staff depending on what hospital you are in, Patient is usually well known, full history unless urgent OT

ALS in ICU

SCENARIO

1. Patient is crashing – 62 year old man been in ICU 1 week, becoming more hypoxic
 - Diagnosis of progressive interstitial lung disease. Lymphoma + other comorbidities
2. Arrest in cardiothoracic ICU – 74 year old man Day 1 post CABG
 - Urgent bedside thoracotomy performed to relieve cardiac tamponade
3. Arrest in a septic patient with severe end stage COPD – limitations of care
 - Sudden LOC, no output

*Notes on ICU – team work, lots of staff usually, Usually not a total surprise
Usually everything is known about the patient*

ALS on the wards

SCENARIO

1. Arrest team – Code blue called to patient in bed space 5, Ward D
 - Code blue called by nursing staff, patient found unresponsive
2. Arrest team – 75 year old lady on the neurology ward, code blue call
 - Team present

*Notes on the ward – Relatively rare, usually lots of staff come running, can be chaotic initially
Usually history well known to their team, often a delay in getting the full story about the patient as team usually not present and nurse may not know all the details of patient. If you are on the code blue team you will know nothing running into the room*

	Scenario #1	Scenario #2
Setting	Busy evening shift in a busy tertiary ED Bed block	
Sx	8yo F BIB mum • Fevers at home, resolved with paracetamol • Mum concerned, child not right • No localising features. No PHx, IUTD • 2 siblings, some contacts unwell with coryza and fevers at school	52 yo F ➢ BIBA decreased LOC (GCS 3), found on park bench ➢ Evidence of head trauma ➢ Likely polypharmacy involved ➢ Patient known to ED and hospital, multiple admissions ➢ Homeless, PHx – Bipolar, T2DM poorly controlled, HT, IHD
Workup	• Exam completely normal • Obs BTF Initially • Increasing RR, Increasing PR, CR prolonged on serial assessment	VBG, US Bloods, CT trauma, Foot Xray
DDx	• FUO - Most likely viral illness. Mother concerned • Obs worsening • Need to consider serious bacterial infection	• Polypharmacy OD – Alcohol, street fentanyl, ICE • Trauma / fall down stairs – Closed head injury, skull fracture ?acute ?old, 1 rib fracture, Ankle closed fracture Weber A • T2DM poorly controlled, BSL elevated 15-20 chronically no acute issues DKA etc
Mx	IVC, IVAB • Admission	• Intubated for control → ICU admission required → BUT still needs to have particular specialty to be admitted under Who to refer? NO other team wants to take her because of her problems: • Neurosurgery - head injury • Gen med (trauma) or gen surg (depends on what is available at hospital) -, HTN, polytrauma • Orthopaedics – if minor – can be managed as outpatient (rib and Ankle ##) • Cardiothoracics • Endocrine consult but NOT admitted under – T2DM review for BSL control • Mental health – polypharmacy, bipolar, homelessness
Patient safety	➢ Patient looks well ➢ Mother is concerned	• ICU admission required → BUT still needs to have particular specialty to be admitted under (but who with) • Ensuring adequate care regardless of social context and previous healthcare experience (beneficence) → recognise the deteriorating patient since high-risk complex patient
Issues	Potential for missed diagnosis and poor patient outcome ➢ <u>Importance of monitoring and vitals</u> ➢ Always look at trends (changes) ➢ Tachypnoea always 1st warning sign <u>Importance of listening to parental concern</u> ➢ They know best Esp for children	• Potential for errors Challenging interactions#1 (if patient has previously assaulted staff member) • Negotiate and de-escalate with nursing or senior staff who are upset • Report any issues to senior consultant (e.g. if senior team talks down on onto junior doctors) – resolve any issues together in small team discussions • Should not jeopardise or delay patient care – the patient still needs to go to ICU (enforce) Challenging interactions#2 (cannot be admitted under any team) • Escalate issue to senior consultant

ACUTE ILLNESS AND UNDIFFERENTIATED ILLNESS


INITIAL APPROACH

- 1) **ABCDE**
- 2) **Why are they here? – DDx** (keep broad – narrow down)
- 3) **Hx- SOCRATES**
- 4) **Systems Review essential** - Headaches → vision → hearing → CADSPIF → Cough, colds, runny nose → N/V, abdo pain, PR bleeding, altered bowel habit → PV bleeding/discharge → muscle weakness/sensation loss → systemic (UWL, fever, NS, rigors, chills) → rashes/lesions
- 5) **PMHx / PSHx / Meds** → Have they been taking **meds = partially treated – ABx?** → meningococcal septicemia
- 6) What does patient and family think?
- 7) "I don't know what exactly could be causing this but we are going to do everything we can to get to the bottom of this!"

Exam	Ix	In ED	ICU	Anaesthetics
1. Vitals 2. CV, 3. Resp 4. GI exam 5. Neurological 6. PR., PV and fundoscopy	Bedside 7. CXR, ECG, UA Bloods 8. FBC, blood film 9. Coags- INR/PT 10. Group and Hold 11. EUC, CMP 12. LFT, BSL 13. CRP 14. Trop/Lipase/BNP 15. B-HCG 16. VBG, ABG 17. Blood culture /swabs Imaging • CT head/chest /AP	18. Have you excluded life and limb threatening diagnosis with appropriate certainty? 19. Can the patient go home or should they be admitted 20. Teams to consult HAVE YOU EXCLUDED LIFE THREATENING ➢ Bleed (perforation) ➢ Infection (sepsis) ➢ Obstruction (bowel, stroke, ACS)	21. Have you excluded life and limb threats 22. Teams to consult 23. Changes to management NEED TO KNOW END DIAGNOSIS	• Can you carry on with surgery • Can the patient continue with prior post op plans or do they need to change ARE THEY FIT FOR SURGERY?

CASE SCENARIOS

	64 year old male presents to ED with Chest pain	6 year old child presents to ED with Rash	64 year old male in ICU day 1 post CABG with chest pain	6 year old child in OT mid elective tonsillectomy with rash
Hx	• 1 day of intermittent chest pain, • exertional, associated with sweating, • He is worried about pneumonia	• 2 days of worsening rash, child otherwise well, ?fever yesterday • Petechial rash, widespread especially lower limbs • Have they been partially treated – ABx?	• Sudden severe chest pain, • getting worse • Feels unwell, • palpitations, • sweating, • no fever	• Urticarial rash, spreading,
Exam	NORMAL – CV, Resp and GI exam	normal	CV and Resp exam Hypotensive, tachycardic, muffled heart sounds, looks unwell.	No airway compromise
Ix	Normal (including CTPA)	Bloods = Plt 8, Other Ix normal	ECG CXR Bloods Bedside echo = tamponade	None
DDx	• ACS • PE, • Aortic dissection • Angina – beware of troponins → crescendo-decrescendo (subcritical stenosis) • AAA • Pericarditis • Pneumonia, pneumothorax • Malignancy • Pleural effusion / pleurisy GORD, referred pain	• ITP • HSP • Meningococcal	Cardiac tamponade	Post-ABx EBV Anaphylaxis allergy
Plan Post Ix	• Admit for Further tests as dx uncertain (e.g. ECG, CXR, d-dimer, imaging) • ECHO – cardiomyopathy • Cardio r/v (or resp)	• Admit • Consult Paeds	• Tamponade needs urgent intervention • Pericardiocentesis	• allergy – needs close monitoring for anaphylaxis



VITALS ARE VITALS

	Reduced	Higher
RR	<u>CNS depression</u> ➤ Centrally acting depressants (opioids, BZDs, alcohol) – overdoses or withdrawals ➤ CNS = Raised ICP, stroke, SCI injury ➤ Electrolyte imbalances ➤ OSA ➤ Hypothyroidism ➤ Sedation	➤ Pulmonary – AOO, asthma, COPD, pneumonia, CHF, PTX, ➤ Cardiac – PE, ischaemia, tamponade, ➤ Haem – anaemia ➤ Endocrine – hyperthyroidism, pheo ➤ Metabolic – acidosis (MUDPILE), severe AKI, CO poisoning ➤ Neuro – post-ictal status epilepticus, brain tumours,
BP	➤ Arrhythmias – bradycardias, tachycardias, fibrillation ➤ Structural HD – valve, IHD, tamponade, HOCM, primary pulmonary HTN ➤ Hypovolaemia – haemorrhage, V + D, diuretics, burns, pancreatitis, diaphoresis (insensible losses) ➤ Systemic vasodilatation – sepsis, anaphylaxis, neurogenic, ANS dysfunction ➤ Obstructive – massive PE, tension PTX, tamponade ➤ Metabolic – hypoadrenalism (Addisonian crisis), hypothyroid ➤ Drugs – BB, CCB, digoxin, opiates, TCA, valproic acid	Non-compliance with anti-HTN + REDCAP ➤ Renal – RAS, nephroblastoma ➤ Endocrine (Conn's, thyrotoxicosis, Cushing's, acromegaly, pheo, congenital adrenal hyperplasia) ➤ Drugs – interactions, OD, withdrawal ➤ Coarctation of aorta ➤ Pregnancy – pre-eclampsia ➤ OSA ➤ Malignant HTN, malignant hyperthermia, Neuroleptic malignant
HR	➤ Physiological – well trained athletes ➤ Hypovolemic shock – haemorrhage ➤ Cardiac = Arrhythmia – Heart block (2nd, 3rd, sick sinus syndrome, CHD) ➤ Infection = endocarditis ➤ Drugs – PSNS stimulation (vasovagal syncope, carotid sinus massage, valsalva) or SNS inhibition (BB, ➤ Endocrine = hypothyroidism ➤ Electrolytes = hyper/hypoK ➤ Hypothermia ➤ Raised ICP (Cushing's reflex)	➤ Pain ➤ Constipation ➤ Exercise ➤ Cardiogenic – Tamponade, ACS, ➤ Resp – PE, PTX ➤ Sepsis, infection ➤ Drugs – SNS (cocaine, ICE) – ingestion, OD, withdrawal ➤ Endocrine = hyperthyroidism, pheo, hyper/hypoglycemia ➤ Anxiety / Panic attack (
Sats	➤ V/Q mismatch – PE, PTX, Pneumonia, APO, ACS, tamponade, hypovolaemic shock ➤ Diffusion issue – APO, pneumonia, ILD ➤ L→R shunting – ➤ Reduced FiO2 – high altitude, ➤ voluntary hypoventilation ➤ other – anaemia, sepsis, foreign body	<u>Always check for (cyanosis with normal Sats)</u> ➤ nail polish ➤ carboxyHb ➤ CO poisoning (smoke inhalation, suicidal attempts) <u>High sats:</u> ➤ Hyperbaric O2 chamber ➤ Hyperventilation – ?Panic attack (Dx of exclusion)
Temp	<u>Exposure</u> ➤ Post-operative ➤ Post-drown ➤ Post burns <u>Late sepsis</u> ➤ Cold sepsis ➤ Hypovolaemic, cardiogenic	<u>Environment / exposure</u> ➤ Athletes / marathon runner ➤ Heat stroke <u>Medical cause</u> ➤ Infection / sepsis ○ Distributive shock – anaphylaxis neurogenic, sepsis ➤ Drug – reaction / withdrawal ○ malignant hyperthermia ➤ Menopause ➤ Vasculitis ➤ Malignancy ➤ Thyrotoxicosis
GCS	<u>BRAIN</u> ➤ stroke – haemorrhagic (EDH, SDH, SAH), clot ➤ SoL – abscess, hydrocephalus, mass ➤ Epilepsy <u>Out of brain</u> ➤ Sepsis ➤ Drug ➤ Electrolytes – low Na, K, PO4, ➤ BSL	<u>Hyperactive</u> ➤ Normal? ➤ Endocrine = hyperthyroidism ➤ Metabolic = ADHD, Mania - bipolar

Things we don't check:

- **Colour** – pale, sallow, jaundiced, well perfused
- **Diaphoresis** – abnormal
- **Posturing** – tripodding, supine, prone, sitting up
- **Exposure** – abdomen, and everywhere else – rashes, bruises, wounds, drains, lines

Making a consult over the phone

- **What is the purpose?**
 - Admission – pick the correct team!!! – escalate to bosses if there is an obstruction
 - Consult
 - Advice
 - Do they need to see/review the patient?
- **Does not necessarily need a full history**
 - E.g. emergency – e.g. a kid that needs tubing OR we need help now!!
- **Aim to be honest, big responsibility**
- **Understand that the receiver will always want to help**

COMMON PRESENTATIONS

	CHEST PAIN	SOB	ABDO PAIN	ALTERED LOC	CRAZY PATIENT
RF	- HC, HTN, DM, - Vascular disease, - inflammatory conditions - Smoking - Drug use (cocaine) - Recent surgery	- HC, HTN, DM, - Vascular disease, - Autoimmune - Smoking - Drug use (marijuana – PTX – THC abuse) - Hx of resp. illness.	- Vascular RF = ischaemic, - Previous bowel surgery - Smoker, EtOH - Autoimmune - Sexually active - Blood thinners (anti-coags)	- Hx of Fits/epilepsy - Prodrome – syncope, chest pain, weakness - T2DM – hypoglycemia - Organ failure - Vascular RF - Drug and alcohol hx - Recent illness	Red flag signs - Known mental health Dx - Drug and alcohol abuse - DM - History of malignancy - Cardio-resp or neuro diagnosis - Older age (> 40yo)
Exam	- Vitals - Sweaty, colour, position, pain - Murmurs - Lung sounds - Calf tenderness (DVT) - Signs of HF (S3, JVP, oedema) Radiation to R) arm more specific for MI	- Vitals - Sweaty, colour, position, pain - Murmurs - Lung sounds - Calf tenderness (DVT) - Signs of HF (S3, JVP, oedema)	Give Analgesia to assist - Vitals - Sweaty, colour, position, pain - CV/Resp exam - Hernia – grey/Cullen, liver stigmata - Genitals – testicles, perineum - Cough/Jump test – ++ peritonism - Palpation – rebound / percussion tenderness - Sign (murphy's, Rosving's) - Peripheries	- Vitals → Postural BP (vasovagal) - Sweaty, colour, position, pain - Exposure → Signs of trauma → sites of infection → rashes - Abdo exam	ENSURE SAFE BEFORE DOING 1st episode of psychosis - Acutely agitated or aggressive patient - Delirium (fluctuating LOC) - Known MH diagnosis attending for review (must medically clear)
Ix	Bedside - ECG – serial - UA Bloods - FBC, EUC, CMP - TFT - Troponin/BNP - D-dimer - CRP - Lipase - B-HCG Imaging - Mobile CXR - Cardiac ECHO - CTPA - CT chest - Doppler LL USS	Bedside - ECG – serial - UA Bloods - FBC, EUC, CMP - Troponin/BNP - D-dimer - CRP (infection) - Lipase - ABG / VBG Imaging - Mobile CXR - Cardiac ECHO - CTPA - CT chest - Doppler LL USS	Bedside - ECG – serial - CXR - Urine B-HCG Bloods - FBC, - EUC, CMP - LFT - CRP - Lipase - Serum B-HCG (ectopic) Imaging - CT AP - FAST / POCUS - US OTHER - Stool culture M/C/S + O/C/P - Urine culture M/C/S - Genital swabs M/C/S	Bedside - ECG – serial Bloods - FBC, WCC - Blood film (schistocytes for TTP) - EUC, CMP - LFT - CRP - VBG – BSL - TFT - Coags - Blood culture - Drug levels – paracetamol, EtOH Imaging - CT brain → MRI, LP - XR – fractures - EEG - POCUS	When to do full work up? 1st psychotic episode - Old pt w/ delirium/ agitation Bedside - Vitals – postural BP - ECG – serial - UA Bloods - FBC, EUC, CMP - LFT, BUN - VBG, CRP, BSL - TFT - B12, Folate, Vit D - ESR - Syphilis, HIV - Drug levels – panadol, EtOH Imaging - CT brain +/- contrast (SOL, bleed) - LP Other - MSE – thought content, suicidal ideation, persecutory delusions - MMSE
DDx	RED FLAGS - ACS - PE - Aortic dissection (diastolic murmur, tearing pain) - Angina – subcritical stenosis (crescendo-decrescendo) SEMI-URGENT - Pericarditis - Tamponade - Pneumonia, - Pneumothorax - Pleural effusion / pleuritis COMMON - GORD - Abdo pathology (referred pain) - Chest wall pain - Costochondritis - rib #	RED FLAGS - ACS - PE - Aortic dissection - Foreign Body - Anaphylaxis - Tension pneumothorax SEMI-URGENT - CCF - Tamponade - AOO – COPD, asthma - APD - Pneumonia - Simple Pneumothorax DO NOT MISS - Anemia - Toxins /ODs (opiates, BZDs, marijuana) - NMD	Bowel - SBO, volvulus - Ischemia, incarceration, strangulation, AAA - Any perforation / rupture (E.g. AAA) - Gastritis, Gastric ulcer, perforated oesophagus - Diverticulitis, appendicitis, abscess - IBD Organ - Hepatitis - GB = stones, cholecystitis, cholangitis, - Pancreatitis, - UTI, Renal colic, stones - Splenic rupture - AAA Gender-specific - Ectopic, endometriosis, adnexal mass/abscess - Ruptured ovarian cyst - Testicular /ovarian torsion Other - AMI - RLL or LLL Pneumonia - DKA - HyperCa	BRAIN ISSUE - Trauma - Vascular – ICH, SAH - Stroke - CNS infection - Raised ICP – SoL (cushing's) - Seizure (post-ictal) DIMEVENT (NON-Brain) - Drugs (OD, withdrawal, interactions – EtOH, BZD, opiates, anti-psychotics) - Infection (UTI, cellulitis, meningitis, encephalitis) - Metabolic (Low Na, low BSL, uremia – AKI, hepatorenal syndrome, UGIB) - Endocrine (Hypoglycemia, hyperglycemia esp. HHS, thyrotoxicosis, myxedema coma, Addisonian crisis, hypoadrenalism – prolonged steroid usage) - Vascular – stroke, HTN, vasculitis, cavernous sinus thrombosis (veins) - Epilepsy – post-ictal, meds, non-convulsive status epilepticus - Nutrition – Vit D, folate, B12, B1 deficiency - Toxins, trauma, TTP	1st presentation – psychosis - Delirium = fluctuating LOC + elderly patient Infection (any sepsis) - AOM, Meningitis - Pneumonia - UTI, cellulitis, pressure sores (nec. Fasciitis) Metabolic - BSL - Electrolytes - Hepatic enceph - Thyrotoxicosis Neurological - CVA - Seizure - Post-ictal seizure - SAH, ICH, SDH - CNS mass Cardiopulmonary - AMI, PE, hypoxia - CO2 narcosis - CCF Drug related - Anti-emetics, anti-histamines, anti-PD - Anti-psychotics - TCA - Withdrawal Other – DO NOT MISS - Pain - Constipation / retention
Risk stratify	- Well's score - PERC for PE	Well's or PERC	Alvarado score	PECARN (for children)	4-AT – orientation, alertness, fluctuating LOC
Mx	Exclude Big 3 – give Mx - ACS = PCI or anti-platelets or anti-coag - Aortic dissection – permissible hypoTN with BB (contact cardiothoracics) If not ACS - Serial ECG and troponin - Observation - Consider D/C 24-hr holter monitor - ECHO - DASS-21 (MH)	ABCDE – ensure airway patent - Avoid lying flat – let patient position themselves - Avoid over-oxygenation in COPD - NIV for COPD and APO Medical - Bronchodilators – wheeze (AOO) 1st line = GTN patch – CCF, APO (or Furosemide 2nd line) - Early ABx – if infective resp. symptoms	- Keep NBM (or last meal) - A – airway patent - B – adequate FIO2 - IVC – bloods - IVF (NS, hartmann) - Analgesia (1g IV/PO Panadol (2/5-5mg IV morphine) (5mg endone PO Intranasal fentanyl - High-dose PPI - D – GCS, BSL Refer accordingly - Gen Surg registrar (urgent if peritonism) - Gastro - Urology - O+G - Vascular	- A – airway patent - B – adequate FIO2 - IVC – bloods - IVF (NS, hartmann) - Support - ABx (if sepsis) - Ionotropes - D – GCS, BSL - Thrombolytics - CVA /interventions - Antidotes for OD (e.g. naloxone, flumazenil) - Give 5mg midazolam for non-convulsive epilepticus to wake them up Refer accordingly - Neurosurg consult - Neurologist - Endocrinologist - Geriatrics	Full work up and investigations? 1st psychotic episode - Old pt w/ delirium/ agitation If known MH dx → medically clear - No further Ix - Await MH review - Document - Avoid premature diagnostic closure General Mx for acutely aggressive pt: - Attempt to de-escalate (offer drink, family support) - Do NOT stand in their way (protect yourself, others & patient) - PO/IM/IV medical sedation (5-10mg diazepam PO = Valium) → required post-monitoring for GCS in resus room - Physical sedation = Security → restrain w/ 5-point immobilization

B – BREATHING & ANAESTHETICS

APPROACH TO RESPIRATORY DISTRESSED PATIENT

Assessment		DDx
Prepare	- Ask (patients, family, paramedics) - Medical records, call GP practice or physician who know patient	Common causes of respiratory distress - Pneumonia - AOO (X-COPD, X-asthma, FB, anaphylaxis, bronchospasm) - APO - PE - Pneumothorax (tension) - Severe hypoxemia Other DDx (ACUTE VS CHRONIC) Heart (tamponade, ACS, pericarditis, LV dysfunction, shunting, arrhythmias) NMD (MG, DMD, paralyzing agent – rocuronium, phrenic nerve palsy) Chest wall (obesity, scoliosis, pectus excavatum) CNS depression (narcotics, stroke, SCI) Poisons / toxins
End-of bed (always look)	- Position (tripod, supine) - Vitals (fever) - Diaphoresis - Pallor	
Respiratory signs (always look)	- O2 requirement - RR - Accessory muscles - Chest expansion - Wheeze, creps	
Systems	- CVS – murmurs, JVP, pulses - Abdo – mass, tenderness, distended	

Risk Stratifying Tool for pneumonia

	Use	Outcome
CURB-65	Predictor of 30 day mortality	- Confusion (AMTS ≤ 8), Urea > 7, RR ≥ 30, BP ($< 90 / < 60$), Age ≥ 65 0-1 (outpatient), 2 (hospital), 3 (ICU)
Pneumonia severity index	Predictor of short-term mortality (higher discriminatory power than CURB-65)	20 variables (demographics, comorbidities, physical exam, vitals) - Investigations also needed (BSL, ABG, CXR)
SMARTCOP	Need for MV or vasopressor	- SBP < 90, Multilobar CXR, Albumin < 35, RR > 25, Tachy > 125 Confusion, Oxygen pO₂ < 70 mm Hg, pH < 7.35, Sats $< 94\%$
PERC rule	Exclude PE	•

INDICATIONS FOR AN ABG!

Why	What	Common Indications
Rapid Results	• Haemoglobin	• Haematemesis • Unexplained haemodynamic instability
	• Sodium • Potassium	• Life threatening hyperkalaemia • Severe hyponatraemia
Assess Respiratory status	• pO ₂	• Technical issues • Abnormal Hb • Severity assessment
	• pCO ₂	• NIV / Intubation • Ventilator adjustment • Extubation
Assess Acid-base status	• HCO ₃ / B.E	• DKA • Renal failure • Overdose
	• pH	• Assess degree of compensation
Specific tests	• Lactate	• Shock • Type B
	• Ionised Calcium	• Massive transfusion • Rhabdomyolysis

DETERMINE DIAGNOSIS – INVESTIGATIONS

Bloods	FBC (Hb, WCC, plt, Hct) EUC (Na, K, eGFR, Cr, Urea) LFT (albumin) Coags D-dimer CRP	Specific Heart Ix ➢ Troponin (beware ONLY raised after 2 hrs) ➢ BNP ➢ ECG ➢ TTE / TOE
Microbio	Blood culture Sputum (? Induced vs tracheal aspirate vs bronchoscopy) Viral swab Urine Beware of contaminants - Coag negative staph in blood - Mouth flora or candida in sputum	Specific Lung Ix ➢ ABG ➢ CXR ➢ US ➢ CT ➢ CTPA ➢ Bronchoscopy
Test	Advantages	Limitations
ABG / VBG	- Rapidly available - Venous ok if pCO₂ normal - Measure of severity "Occult" hypercarbia	- Does not frequently guide initial management - Painful / Stressful for patient
CXR	- Rapidly available - Cornerstone for diagnosis (must compared with previous CXR)	Sensitivity and specificity Inter-interpret variability
CT	- Pulmonary Emboli - Cancer - More sensitive and specific for other causes	- Patient needs to lie flat (bad if haem compromised) - Contrast allergy or impaired AKI (DM, CKD) - Radiation load - Resource burden
US	- Rapidly available - Concurrent assessment of heart and lungs - Broad division (Airways disease, Interstitial, Pleural)	- Appropriate training required - Incorrect interpretation
Bronchoscopy	- More sensitive for less common pathology - More likely to tailor ABx - Possible therapeutic benefit	- Well enough to tolerate or be intubated - Complications (hypoxia, aspiration, pneumothorax)

RESPIRATORY CARE MX – (1) Supportive Care + RESUS → (2) ABCDEFG → (3) Therapy Goals

1) Supportive care and Rx symptoms (SOB, chest pain, vomiting) → FiO₂, analgesia, anti-emetics 2) Ensure adequate oxygenation & ventilation (CO₂ elimination) - What's the goal? - PaCO₂, pH or clinical end-point - Methods improving ventilation? → control FiO₂, NIPPV, intubation - Correct reversible pathology – bronchospasm, sedative toxicity 3) Optimise haemodynamics + fluid status - HTN (High LV afterload – APO) → SL/IV GTN - HypoTN (?shock → hypovolaemic, obstructive) → IVF - Arrhythmias (Atrial tachycardia esp. in acute resp. failure) → rate and rhythm control	Goals of therapy 1) Burden on patients/family - uncomfortable support NOT treatment - prevents patients from communicating 2) Prognosis in those with resp. failure with advanced disease – - burden of care, - likelihood of survival - perceived QoL
--	--

	Minimally invasive	Flow Rate	FiO ₂ (%)	Indication for use			
	Nasal Cannula/Prongs	1-4 L/min (turned up in ED)	25-35	- Minimal resp. distress (≈90%) Inverse rule (FiO₂ ≈ 1/flow) → Lower FiO₂ for greater WOB - Less claustrophobic, low cost, can eat and speak			
	Simple Face Mask	5-10 L/min	40-60	- Moderately hypoxic - Risk of CO₂ retention			
	Venturi Mask blue – white- orange – yellow – red-green (2 – 4 – 6 – 8 – 10 – 15)	2-10 L/min (adaptor)	24-60% (fixed)	- COPD patients (i.e. avoid CO₂ retention) = loss of hypoxic drive to breathe - Transition to non-rebreathing mask in emergency or if patient is not well			
	Non-Rebreathing Mask "Reservoir bag MUST be inflated + tight fit needed"	15 L/min	> 60%	- Oxygen reservoir to give higher FiO₂ with adequate ventilation Mainly for - Post-cardiac or respiratory arrest, - Severely hypoxic patient			
	Hi Flow Nasal Cannula "humidified O ₂ delivery under pressure"	≤60 L/min	Up to 100% (hard to titrate)	- Low level prep → applied nasally or via tracheostomy - Humidifier replicates humidifying effect of nose Matches inspiratory flow rate even with increased WOB - Aims to reduce subjective WOB and SOB			
	Bag-Valve- Mask Resuscitator	15 L/min	>0.8	Key principles: Almost always Need adjunct → nasopharyngeal or oropharyngeal (Guedel) airway or BOTH - optimize patient position (inc. bed height) Difficult BVM causes - BONES - Beard = impedes a good seal - Obese - No teeth = loss of facial architecture – (but easier intubation) - Elderly - Sleep Apnea / Snoring			
	Non-invasive PPV (CPAP, BPAP)	Adjustable L/min	Fixed FiO ₂ (up to 100%)	CPAP Constant pressure → alveolar recruit - PREVENT airway collapse ↑↑ atelectasis + alveolar Vent. - Reduces LV afterload	T1RF	BIPAP IPAP (air forced into lungs) + expiratory PEEP (prevent collapse) - Reduces inspiratory effort - Increased tidal volume and CO₂ clearance	Relative CI - Facial trauma - Recent surgery (URTI, upper GIT) - Reduced LOC - Aspiration risk (obtunded pt) - Pneumothorax - Signs of RV strain
	Mechanical ventilation OR Tracheostomy (emergency)	Adjustable Control FiO ₂ , RR, TV, Peak flow rate, PEEP	100%	Endo-tracheal tube = definitive airway to maintain airway patency LMA (laryngeal mask airway) = non-definitive airway Dynamic manoeuvres (e.g. prone position) - reduces lung compression by adjacent organs - improves ventilation and perfusion PLUS improves oxygenation and secretion clearance			

***** **Fraction of Inspired Oxygen (FiO₂) concentration ≈ (Flow Rate x 4) + 21 (FiO₂ in room air)** *****

	WHAT IS IT?	INDICATIONS	SIDE EFFECTS
Anti-microbials	Infection	Organisms	Anti-microbial
	CAP	Streptococcus, Haemophilus, Moraxella	Ceftriaxone
	Atypical	Mycoplasma, Legionella, Chlamydia	Azithromycin
	Hospital Acquired	- Resistant Gram-ves (pseudomonas) - MSSA vs MRSA	Tazocin Vancomycin
	Chronic lung disease	Pseudomonas	Tazocin
	Immunocompromised	- Neutropenic - PJP - Fungal (e.g. aspergillosis) - Viral reactivation (CMV)	Tazocin & gentamicin Bactrim Antifungal Valganciclovir
	Viral	Influenza	Oseltamivir
	Abscess	Usual + Anaerobes	Tazocin
Bronchodilators	INHALED	- Neb SABA (salbutamol 5mg) - Neb SAMA (ipratropium 0.5mg)	- Viral induced wheeze - Acute exacerbation of AOO (asthma, COPD)
	SYSTEMIC	- IV Salbutamol (only if PO not tolerated) - IV/IM adrenaline - IV MgSO₄ - Aminophylline - Ketamine	- Anaphylaxis - SMC relaxation for severe exacerbations - Intubation or to tolerate NIV
Chest drains	- Needle - 2nd ICS, MCL - Finger thoracostomy - Aspiration ONLY - Pleural caths	- Tension pneumothorax → URGENT decompression 2nd ICS, MCL - Pleural effusion (exudate, transudate, malignancy, empyema) - SOB/hypoxemia - Haemothorax → REQUIRES a drain	SABA – Tachycardia, tachyphylaxis, lactemia Ipratropium – antichol / (anxiety, tachycardia, dry secretions) MgSO₄ / aminophylline = flushing, hypoTN
Diuresis	- Furosemide 20-80 mg IV "Dry lungs are happy lungs"	- APO - Cor pulmonale	- Re-expansion pulmonary oedema - Pneumothorax - Infection (pleuritis) - Adjacent organ damage (liver, RA)
Expectoration	- Chest Physiotherapy - Mucolytics - Humidification via HFNO - Bronchoscopy	- CF / Bronchiectasis - Pneumonia - well enough to tolerate or intubated	
Fibrinolysis	SHOULD DO CTPA 1st is possible!!! - Tissue plasminogen activator (TPA) - Alteplase / Tenecteplase	Confirmed massive PE (Haemodynamic instability) Probable massive PE too unstable for imaging Submassive PE (Right heart strain) Syncope (stroke, ACS) → CT brain to exclude ICH prior	Bleeding risk – internal bleed, or haemorrhagic stroke
Glucocorticoids	Oral pred or IV hydrocortisone, O/IV dexamethasone - If tolerating oral meds and gut working oral and IV have <u>equivalent</u> efficacy	Exacerbation of asthma/COPD → reduce hospital stay Anaphylaxis - Exacerbation of steroid responsive lung disease (Fibrosis, Allergic Alveolitis, Pulmonary Vasculitis +/- autoimmune) - Consider in COVID-19 and CAP →	Acute complications of steroids: - HIGH BSL - Fatigue / weak - Immunosuppressed - delirium/psychosis

ABG INTERPRETATION

How to interpret ABG/VBG?

- Check pt details and ALWAYS CHECK OTHER RESULTS (Hb, lactate, BSL, K)
 - Circle HIGH and LOW values
 - Is it venous or arterial? → Do I need ABG or VBG?
 - If ventilating/sepsis = **need ABG**
 - If chronic lung disease/DKA = **VBG + SaO₂**
 - DO NOT DO ABG ON ROOM AIR IF PATIENT IS HYPOXIC**
 - If **resp. acidosis** → check **A-a gradient** (diffusion issue?)
 - High A-a = lung issue (NOT CNS)
 - If **NO resp. or metabolic cause** → check **Base Excess** (-2.5 to +2.5)
 - < -2 = acidosis
 - > 2 = alkalosis
 - Is there another cause of the acidemia or alkalemia?
 - If **metabolic acidosis** → check **Anion Gap** to determine if:
 - (1) **intrinsic loss** (i.e. renal, DKA, Lactate) or
 - (2) **extrinsic acid** (e.g. methanol, anti-freeze)
 - Respect buffering** (HCO₃ may be normal) → ECF (HCO₃) Blood (Hb), ICF (PO₄) buffering
 - Summarise findings**
- This patient is a _____ aemic Due to a primary _____ osis With partial/full _____ compensation



Severity of T1RF: P:F ratio or S:F ratio [ONLY from ABG]

$$P:F \text{ ratio} = \frac{PaO_2}{FiO_2} \text{ or } \frac{satsO_2}{FiO_2}$$

In ARDS = <300 (mild), <200 (moderate), < 100 (severe)

A-a gradient (difference b/w O₂ conc. in alveoli and arterial system)

$$A-a \text{ gradient} = P_{A_{O_2}} - PaO_2 \approx \frac{Age}{4} + 4$$

$$P_{A_{O_2}} = FiO_2 \times (P_B - P_{H_2O}) - \frac{PaCO_2}{RQ}$$

Cause for hypoxia	A-a gradient	Corrected with high FIO ₂ ?	Causes
Low FIO ₂	Normal	Yes	High altitude, hypoxic gas mixture
Hypoventilation	Normal	Yes	Residual anesthetic, muscle relaxants
Diffusion	Elevated	Yes	Interstitial lung disease
V/Q mismatch	Elevated	Yes	Mucus plug, pulm embolism, COPD
Shunt	Elevated	No	Atelectasis, ARDS

First formula:

$$\text{Anion gap} = Na^+ + K^+ - (Cl^- + HCO_3^-)$$

Value of Anion Gap

normal:

8 - 12 mEq/L (if without potassium)

Second formula:

$$\text{Anion gap} = Na^+ - (Cl^- + HCO_3^-)$$

12 - 16 mEq/L (if potassium is given)


Is there an anion that is NOT measured? E.g. albumin



Metabolic acidosis = uses up HCO₃ → Low HCO₃ on ABG/VBG

	pH	pCO ₂	HCO ₃ ⁻	Cause	
Normal	7.35-7.45	35-45mmHg	22-26 mmHg		
Resp. acidosis			Normal (ACUTE) *mainly from intracellular Hb, PO ₄ buffering		
			↑ (CHRONIC) * ↑↑ 1mM HCO ₃ = ↑↑ 10mmHg PaCO ₂ (occurs after 24 hrs w/ max response at 3-4 days)	Acute - Fever - Increased catabolism - Sepsis - Steroids - Burns - Pancreatitis - Thyrotoxicosis	↓V_A (↓Alv ventilation) AP₀, pneumonia (T2RF + raised A-a) CNS depression = Narcotics / opiates, Post-op Sedatives, stroke, SCI Upper airway obstruction (aspiration, laryngospasm after intubation, anaphylaxis) Hypoventilation (T2RF + normal A-a) Mechanical ventilation (hypoventilation, permissive hypercapnia)
Resp. alkalosis			Normal (ACUTE) *mainly from intracellular Hb, PO ₄ buffering (within 10mins - max response at 6 hrs)	CNS	- Pain - Agitation / Panic = ↑PO₂ - Encephalopathy (uremia, hepatic, infective)
			↓ (chronic) ↓ 5mM HCO ₃ = ↓ 10mmHg PaCO ₂ (from 6 hrs to 2-3 days)	Lung PE, pneumothorax → ↓PO₂ XS ventilation (mechanical) → good V, poor Q High altitude - Mt Everest Compensatory Kussmaul - DKA Other VERY ILL = Sepsis!!! - Drugs - salicylates - Progesterone	Upper airway obstruction (OSA) Chest wall deformities (scoliosis, pectus excavatum, rib fracture) X-Lung disease (COPD, asthma) NMD (MG, DMD, neuropathy) Obesity hypoventilation syndrome
Metabolic Acidosis (L TKR)				High anion gap metabolic acidosis (HAGMA) Toxins → MUDPILES Lactate - IV adrenaline - Metformin Urate or XS H⁺ (RENAL FAILURE) - XS PO₄, sulphate, Hippurate - Type 1 RTA - rhabdomyolysis Ketones - DKA (T1DM, SU, Insulin) - Starvation state (AN) - Alcoholic Alcohol (toxins) - Ethylene glycol (anti-freeze) = osmolar gap - Pyroglutamic acid = XS N-acetyl cysteine	
				Normal anion gap metabolic acidosis (NAGMA) Increase Cl (negative UAG) - NaCl - TPN GI HCO₃ loss (negative UAG) - Small bowel (diarrhoea) - Stoma output - Pancreatic / biliary loss / drains Renal HCO₃ loss (positive UAG) - RTA - Acetazolamide - HypoAldo (spironolactone) - compensation	
Metabolic alkalosis				Above Stomach loss = VOMITING, NGT/NPA aspiration Exogenous HCO₃⁻ = Overcorrection IV or Milk alkali syndrome (XS Mylanta - antacids) K⁺ depletion → Cushing, diuretic +/- diarrhoea HyperALdo → Conn's, cirrhosis, HF, loop and thiazides	
Mixed acid base	- Mixed metabolic acidosis/alkalosis - Mixed metabolic and respiratory			- Lactic acidosis/DKA with vomiting (met acid + alkalosis) - Salicylate poisoning (met acid + resp. alkalosis)	

CXR Checklist [PA = good, AP = crAP!]



First pass	Review areas
Trachea	
Mediastinum	
Cardiac borders	Pneumomediastinum
Costophrenic angles	Retrocardiac
Soft tissues	Apices
Bones	Below diaphragm
Below diaphragms	

1. Intro (Pt Name, DOB, date, projection, quality → RIPE)

- Rotation** – medial border of clavicle heads equidistant to spinous process
- Inspiration** – see 6 anterior ribs, MCL
- Position** – supine, erect, lateral, decubitus
- Over-exposure (black)**, under exposure (white) – VERTEBRAE JUST VISIBLE BEHIND HEART

2. AIRWAY- tracheal/mediastinal deviation (? patient rotation) OR obvious masses

3. BREATHING – lung fields, pleura, hilar region

- Air** (?pneumothorax, emphysema)
- Fluid /meniscus sign**(? Effusion)
- Consolidation** (?infection)
- Lobar collapse**
- Lesions** (e.g. malignancy, abscess)

Pleura

- Pleural plaques (?asbestosis)
- Calcification

Hilum

- Bilateral enlargement (?sarcoid)

4. BONES/BREAST SHADOW

- Obvious fractures- clavicles, flail ribs
- Scoliosis, lytic lesions
- METs
- Pectus excavatum

5. CIRCULATION


- Cardiomegaly 2° to HF** (> 50% cardiothoracic ratio – adults, > 65% infants)
- Heart border (R = RA, L = LV)
- Aortic knuckle should be visible
- Mediastinal width (>8cm = **aortic dissection**)

6. DIAPHRAGM

- high riding** – phrenic nerve palsy, DMD, MND, MG
- Flattening** – obstructive lung disease (emphysema, asthma)
- Air under diaphragm – pneumoperitoneum (abdo perforation)
- Blunting costophrenic angles – pleural effusion

7. EXTRA DEVICES

- CENTRAL LINE** = internal jugular vein → CVS and cavo-atrial junction
- NGT** =below diaphragm at tip of stomach
- PICC line** = peripheral → cavo-atrial junction
- Portocath** = Subclavian vein → cavo-atrial junction
- ETT** = mouth →3-5cm above carina (level of aortic knuckle)
- Chest drain** = in pleural space
- ECG electrodes**
- Sternal wires** (previous open heart surgery)



Key knowledge:

- BLACK**: air > fat > soft tissue > bone **WHITE**
- Describing abnormality:**
 - Density** (relative to what?, uniform vs patchy?)
 - Left vs Right** (Which zone?)
 - Anatomical position** (pleura vs parenchyma)
 - Size + borders**
- MOBILE CXR** → **LARGE HEART + SEVERE ROTATION**

Summarise


- This is an erect PA CXR of ____ showing _____. To complete my analysis, I would examine previous films and ascertain the clinical hx
- My ddx _____
- Investigations I would order _____

Common presentations/DDx

Opacification (NOT just infection)	Diffuse alveolar shadowing
- Non-uniform opacification indicating substance in alveoli air-bronchograms (visible bronchioles penetrating consolidated area)	Alveolar oedema caused by fluid filling the alveoli and small airways may typically see "bat wing's distribution"
DDx: for opacification	Basic
Fluid (pus, serous, blood) Infection / Pneumonia Atelectasis / collapse Mass / Cancer Trauma / contusion Infarction Emboli	- Pneumonia (viral – COVID atypical, pHP) - APO - Pulm. Fibrosis
	Extended
	- Pulm. Haem - Extrinsic allergic alveolitis - Vasculitis - Lymphangitis - Miliary pattern (TB, old chicken pox)














Opacity	Pathology	Opacity (L)	Pathology
R paratracheal stripe	RUL Ant. Mediastinum	Aortic knuckle	LUL
R heart border	RML	L heart border	Lingula
R hemidiaphragm	RLL	Left hemidiaphragm	LLL

*Nb: Can have mixed pathology (both lobes affected)



Pneumonectomy	- No lung markings - Mediastinal and tracheal shift
Pneumoperitoneum	- Perforated viscous - Post-op laparotomy
WHITE-OUT	White out of mediastinum: <u>Towards</u> = collapse <u>Away</u> = effusion <u>Middle</u> = both Other causes = ARDS - Sepsis, - trauma, - DIC, - burn, - O2 toxicity, - fat embolism

COMMON PRESENTATIONS

			
Pneumonia	Infarction	Trauma	Cancer
			
Haemorrhage	Atelectasis - Affected lobe is smaller - Raised diaphragm ipsilateral - Structures moved <u>towards collapse</u> <i>Cause: obstructive, pneumothorax</i>	Emboli	APO Alveolar shadowing (bat wing) - Kerley B lines (i.e. <i>lymphatic congestion in lateral lower edge</i>) - Kerley A - apical - Kerley C - central Cardiomegaly - Upper lobe venous Diversion Effusion Rx: - GTN - NIV ventilation - Diuresis w/ furosemide (if BP stable)
			
Pneumothorax - Best on expiration film - Absent lung markings - Mediastinal shift - Surgical emphysema?	Tension pneumothorax Cannula 2nd IC MCL Finger thoracostomy Chest drain	➤ Surgical emphysema	Effusion vs mass ➤
			
Supine pneumothorax - Deep sulcus sign - Lung edge - Mach effect	L pleural effusion - Dense opacification + Blunting CP angles + Meniscus sign Transudate (nephrotic, liver, CCF e.g. dilated CM, thyroid, ovarian - Meig's, enteropathy) Exudate (PE, infection, infarction, inflammation, neoplasm, TB) General Rx ➤ Pleural tap – send for cytology, LDH, pH, protein, glucose ➤ Chest drain	Lobar Collapse ➤ Endoluminal (sputum, blood) ➤ Luminal (cancer, infection e.g. acute bronchitis) ➤ Extraluminal (LN, cancer) General Rx ➤ Early Abx (e.g. augmentin) ➤ Chest PT ➤ CT to exclude malignancy	Focal infiltrate in HTN and smoker ➤ Pneumonia AND ➤ Cancer General Rx ➤ Early ABx for CAP (7-10 day benPen 1g tds PO) ➤ Repeat CXR to confirm resolution

C- CIRCULATION / SHOCK

Diagnosis of shock

Shock is a cellular state of hypoperfusion

Hx	• Collapse, syncope • Signs of End-organ hypoperfusion (chest pain, confusion, abdo pain, oliguria) • Compensatory signs – WoB, kussmaul, polydipsia	
Exam	• Witals → ↑RR, ↑HR, ↓BP, Sats, Temp • Warm - Capillary refill time, pallor, diaphoresis • Wet /dry (fluid status) – oliguria, MM, bibasal crackles, oedema • With it - GCS/Confusion / Drowsiness	
Tests	• Wactate - Lactate (VBG) – good to monitor to response of treatments • CRP	
Non-invasive tests	Semi-invasive tests	Invasive tests
• Existing monitoring – Sats, ECG • CXR • TTE • Impedance	• Arterial Line • TOE • Oesophageal doppler	• PICCO – trans-pulmonary thermodilution • Swann-Ganz – right heart thermodilution • CVO₂/MVO₂


General Acute Mx of shock

- 1) Triage - CAT1-5
- 2) Transfer to appropriate area (e.g resus, bed)

A	Unconscious / Procedures
B	High work of breathing → sats, RR, FiO ₂
C	Arrhythmias / Cold or warm peripheries ➢ ECG ➢ IV Access x2 → Bloods – FBC, EUC, LFT, troponin, VBG, culture M/C/S ➢ USS – FAST ➢ UA Optimize hemodynamics 1. Check if fluid responsive 1st a. Straight leg raise, calf pump or lie supine to increase VR 2. Restore intravascular volume "like for like" a. IVF crystalloids vs colloids (alternate each day – reduce acidosis) b. Listen to chest for APO – <u>bibasal crackles</u> → consider reducing fluids and adding FiO₂ 3. Maintain BP with vasopressors MAINLY → target MAP > 65, SBP > 90 4. Assess and optimize CO 5. Decide when to stop (e.g. not working or APO) → DO CXR!!
D	Delirium – 4AT
E	Bleeding, Melaena, Hypothermia – check for rashes
G	Hypoglycemia, abdo exam,

Types of shock + Mx

SIRS	<u>(at least 2 of these to meet criteria)</u> 1. Body temp >38°C or <36°C, 2. HR > 90/min 3. RR > 20/min or PaCO₂<32mmHg 4. WCC >12.0 x 10⁹/L or <4.0 x 10⁹/L	<u>Pathophysiology:</u> 1) Raised pro-inflammatory mediators (IL-1,6 and TNF-α from macrophages activates neutrophils, platelets and endothelial cells) → DIC + increased vascular permeability 2) Raised acute phase proteins (buy time) 3) HypoTN, hypoperfusion, hypoxia
Sepsis	Life-threatening organ dysfunction caused by a dysregulated host response to infection	
Severe Sepsis	Sepsis + evidence of organ dysfunction (e.g. hypoxia, oliguria, AKI, Coag dysfn, hypoTN, raised lactate > 2mM, thrombocytopenia)	
Septic Shock	- Sepsis + arterial BP drop causing organ hypo-perfusion ➢ Systolic BP < 90 (DESPITE adequate fluid resus) ➢ Raised lactate > 4mM	
Septicaemia	Bacteria enters bloodstream triggering sepsis (e.g. meningococemia)	
Bacteraemia	Bacteria in the bloodstream. May or may not cause SIRS or sepsis.	
Risk factors	- Extremes of age (<1 yo or > 75 yo) - Chronic conditions (e.g. COPD and diabetes) - Chemo, immunosuppressants and steroids - Recent surgery, trauma or burns - Pregnancy or peripartum - Indwelling devices (e.g. catheters and central lines)	




qSOFA score

IF SCORE > 1 THEN INVESTIGATE FOR PRESENCE OF ORGAN DYSFUNCTION AND INCREASE FREQUENCY OF MONITORING

Sequential Organ Failure Assessment → measure mortality in ICU

- Quick SOFA (≥ 2 of the following = **HIGH** Mortality)
- GCS < 15, RR ≥ 22, sBP < 100
- Febrile
- Reduced UO- cloudy urine

Types of shock + Mx

Shock type	Hypovolaemic	Cardiogenic	Distributive			Obstructive
			Neurogenic	Septic	Anaphylactic	
Cause	Haemorrhagic (big 5 → chest, abdo, external, long bone, pelvis) > Aortic dissection > UGIB/LGIB > Mallory Weiss / Boorheave > menorrhagia Non-haemorrhagic - Electrolyte imbalance (V + D, Dehydration, insensible losses, pancreatitis) - Burns - Inadequate intake (long lie / immobility)	Electrical - Tachyarrhythmia - AF + RVR, VT/VF (AF → ↓ EDV → ↓ CO by 20%) - Bradycardia - complete HB, BBB - Drugs = BB, CaB, digoxin, sotalol, amiodarone Structural - Myocardium = Dressler, CHF, CMP free wall rupture, myocarditis - Valvular = AS, endocarditis (IE), AR, valve rupture, papillary muscle rupture	- paraplegia, - acute SCI - spinal anaesthesia SNS inactive – no signals sent to brain → vasodilation of SVR	40% mortality rate - Bacteria - Gram +ve/-ve - Fungi - Virus - Protozoa - Causes pain (if attacks nerves) Other distributive causes: - Endo: Adrenal / thyroid / vasopressin insufficiency - Electrolytes: CMP, EUC - Toxins – alcohol - Drugs – Metformin, serotonin syndrome, aspirin	- Drugs (ACEi, C1 esterase def., anaesthetics) - Food / bites - Idiopathic DDx (SCRAM) - Scrombotoxin - Carcinoid - Red-man syndrome - Alcohol-related - Medullary carcinoma of thyroid - Acute GORD - Dystonic reaction	↓preload + ↑afterload (↓venous return) - Tension pneumothorax - Massive PE - Cardiac tamponade (blood, serous fluid) - Pleural effusion - Air embolism
CR	Long + Cool/dry		Short + Warm/dry (due to vasodilation)			Long + Cool/dry
HR	↑↑↑	↓↓↓	↓↓↓	↑↑↑	↑↑↑	↑↑↑
CO:SVR	CO < SVR	CO < SVR	CO > SVR "heart squeezing hard but high distribution/leaky BVs makes it difficult to perfuse organs"			CO < SVR
Ix	- FBC (blood loss) - EUC - CXR - Pelvic XR (pelvic #) - eFAST (abdo)	Examination - murmurs ECG signs: - AMI signs, LVH, SVT CXR = Acute pulmonary oedema ECHO – determine need for fluids vs inotropes	- Sphincter dysfn - Paraesthesia - Weakness - Signs opposite to typical shock	3 tests: - Blood lactate (VBG) 2x sets Blood cultures - Urine output 3 treatments: - O2 (maintain 94-98%) - Empirical broad spectrum Abx - IVF +/- inotropes (maintain MAP >65 mmHg)	- Rash (urticaria) - Wheeze - Gut pain, Oedema, Angioedema – IgE released → mast cell degranulate → histamine → vasodilation -	Tension PTX - Distended neck veins - Dev. trachea Beck's Δ (tamponade) - HypoTN - Muffled heart sounds - Distended veins
Rx/Mx	Acute Mx: - Warm IV bolus NS (500mL in < 15 mins) – lower if elderly - Vasopressors (↑ SVR) - TXA - Group + X-match (pRBC with O neg) Surgical Mx: - Surgical /radiological / endoscopic surgery → trauma, blood loss, intra-abdominal issue - Embolise bleeding vessel - Endoscopic clipping (variceal bleeds) Maximizing oxygen delivery → maintain tissue perfusion - Target MAP >65 - normalise lactate - CVP monitoring - Target CPP = 50–70 mm Hg (as per BTF guidelines) - Intubation, ventilator (if ≤30 cmH₂O) - Transfusion (if Hb < 70, plt < 20) High vol. hemofiltration in severe met acidosis or renal failure	- Inotropes MAINLY (e.g. dobutamine, adrenaline) to increase cardiac output 'Fluid challenge rarely useful (NOT fluid responsive)' To improve diastolic filling of coronary arteries and reduce afterload - Mild hypothermia can improve neuro outcome of STEMI - CT Angiography - PCI - stents - Thrombolysis  'intra-aortic balloon' - TOO HIGH = block L SCA - TOO LOW = block renal flow (pre-renal)	Check for bleeding 1 st (? spinal trauma) e.g. may be bleeding out but is on a BB - Rule OUT hypovolaemic shock - Rule OUT drugs and poisons (e.g. opioid, nitrate, BB) - Avoid over rehydration with fluids - Beware that vasopressors reduce BP	Hemodynamic resus (SEPSIS 6) - Treat within 1 hour - IV fluids crystalloids 20mL/kg - NS vs Hartmann - Monitor acid-base - Vasopressors 1st line: NORAD, metaraminol, vasopressin - adrenaline (for arrhythmias or splanchnic ischaemia) Early Abx (within 1 hr after cultures) - Local guidelines – vancomycin, tacomycin → narrow spectrum (after culture result) - Source control = debride, remove, drain septic source, repair perforation Maximizing oxygen delivery → maintain tissue perfusion - Target MAP >65 - normalise lactate - CVP monitoring - Target CPP = 50–70 mm Hg (as per BTF guidelines) - Intubation, ventilator (if ≤30 cmH₂O) - Transfusion (if Hb < 70, plt < 20) - High vol. hemofiltration in severe met acidosis or renal failure	- Stop source/allergen - Lie flat - IM adrenaline (500mcg – adults) (300mcg – >6 yo, pregnant) (150mcg < 6yo) - High flow FIO₂ - Monitor IV fluid (if hypoTN) - IV/IM hydrocortisone 200mg of 4mg/kg For children CPR: 1-hand (≤ 8) 2-hand (> 9) - Chest compress (100-120bpm) 15 comp/2 breaths for all children	Bedside - ECG - Right heart strain (for massive PE) – T wave inversion in INFERIOR LEADS AND V1-3) - CXR = Pneumothorax - PE = thrombolysis (TPA agents) +/- embolectomy - PTX = needle decompression - Tamponade = pericardiocentesis

5) Cervical shock – Missed M/C → HypoTN in high -B-HCG women → need to remove POC (Avoid XS fluid resus)

6) High Output Heart Failure shock (XS Vasodilation + XS O2 demand → chronic ↑CO → chronic ↑preload + ↓SVR)

- beri-beri (systemic vasodilation),
- thyroid storm (thyrotoxicosis),

- obesity
- Myeloproliferative disorder

- severe, anaemia
- Paget's disease

- Cirrhosis
- AV fistulas

Medications to optimize MAP = Guarding the Perfusion Pressure

General Principles


- Ensure adequate organ perfusion (MAP) to prevent end-organ dysfunction (MOD and death)

Don't forget to treat the underlying cause e.g.

- **Fluids** → hypovolaemic shock
- **Thrombolysis** (streptokinase) → massive PE
- **PCI** → cardiogenic shock due to MI
- **Adrenaline** → anaphylaxis

Use vasoactive agents to maintain Blood Pressure

- Typically Mean Arterial Pressure (MAP) > 65 mmHg)
- Metaraminol in **boluses or infusion**
- Noradrenaline or adrenaline as **infusions**




* **Excessive** increases in either CO or SVR does **NOT** restore normal organ perfusion despite adequate blood pressure

	MoA / drug class	HR (chronotropy)	SV (inotrope)	SVR
Positive chronotropy				
Isoprenaline (synthetic)	B-agonist			
Atropine, glycopyrrolate	Anti-cholinergic			
Negative chronotropy				
Labetalol	NSBB + alpha blocker			
Diltiazem	CCB			
Digoxin, amiodarone	Anti-arrhythmias			
Negative chronotropy + inotropy				
Metoprolol	Cardio-Selective BB			
Verapamil	CCB – avoid in HF or LVF			
Positive inotropy				
Dobutamine (Catecholamine)	B agonist + chronotropic +)			vasodilator
Milrinone (PD3E inhibitor)	intracellular Ca ²⁺ + some chronotropic			
Levosimendan (PD3E inhibitor)				
Positive inotropy + Vasopressor				
Adrenaline (Catecholamine)	B agonist (low dose), A – agonist (high dose) Anaphylaxis, hypoTN			High dose
Vasopressor – alpha agonists				
Noradrenaline (vasopressor)	a1 agonist - ONLY via central line IV		High dose – due to high SVR increasing LV afterload to reduce SV	
Metaraminol (vasopressor)	a1 agonist - peripheral IV or bolus – causes reflex bradycardia and tachyphylaxis)			
Vasodilators				
GTN, hydralazine	Nitric oxide donors			
Amlodipine, Clivedpine	CCB			
Prazosin	a-Blocker (beware of postural HypoTN and reflex tachycardia)			
FINAL RESORT				
Sedatives	Sympatholysis – bradycardia and vasodilaiton			

CARDIAC MONITORING

What?	Why?	How?
ECG - Rate - Rhythm - ST segments - PR / QT intervals	- Arrhythmias - Ischaemia - Drug toxicity - Electrolytes	- Repeat 12 lead - Continuous
Blood pressure - SBP - DBP - MAP - Pulse pressure - Variations with respiration	- Vasopressor requirement • Risk of deterioration	- NIVBP - Arterial line
Cardiac Output - Cardiac Index ?volume responsiveness ?volume overload	- Assessment - Response to therapy - Guide therapy	- ECHO - Pulse Contour - Thermodilution

ARTERIAL LINE




	Advantages	Limitations
Non-invasive	- Instantly available - Avoids complications of arterial line	- Limited time cycle (i.e 1 per minute) - If required frequently - Irritating (sleep etc) • Bruising - Cannot be done over PICC lines
Arterial line	- Continuous monitoring • Phlebotomy - Monitor lactate - Additional information - Cardiac Output - Respiratory variation	Complications - Insertion (arterial injury) - Distal ischaemia - Exsanguination - SBP and DBP may be unreliable - Limit patient mobility

CARDIAC OUTPUT MONITORING

Procedure	Waveform
Thermodilution CO Cold saline injected into central vein - Measures temp. change in antegrade BV - Pulm. Artery (R heart) - Femoral artery - Area under curve = inversely proportional to CO	
Pulse Contour Analysis - PiCCO Uses femoral artery line and upper body CVC - Check the trends	
Pulmonary artery catheter - Gold standard of CO monitoring - Uses pressure transducer and thermistor in pulmonary artery <u>Complications</u> ➢ Ruptured pulmonary artery ➢ VT	

GENERAL ARRYTHMIA MX:



PC: 65 M w/ 3 hrs of palpitations, light-headedness, chest pain appearing pale and diaphoretic


Working out

1. Fast AF
2. Inferior ST elevation
3. Anterolateral ST depression

Dx = myocardial ischaemia with unstable rapid AF

Mx:

1. Rate control – BB, CaB
2. Rhythm control
3. ACS treatment – PCI → BB, aspirin, statin, clexane



PC: 42 F w/ altered mental state and PMHx of epilepsy presents with bradycardia and hypotension


Working out

1. Complete heart block
2. Inferior ST elevation
3. Lateral ST depression

Dx = inferior MI associated with complete heart block

Mx:

1. Speed up rhythm
2. PCI



PC: 85 M w/ coffee ground vomit since AM and severe epigastric pain


Working out

1. Sinus rhythm
2. ST elevation in AVR
3. Global ST depression

Dx = diffuse endocardial ischaemia secondary to UGIB

Mx:

1. Wide bore IV access
2. Check Hb and transfuse > 80
3. PPI and endoscopy



PC: 55 F called Ambo due to chest pain. Now confused and unable to give history


Working out

1. Complete heart block
2. Left axis and LBBB morphology
3. Abnormal ST depression V2/3

Dx = Acute MI

Mx:

1. ACS protocol
2. PCI



PC: 25F found unconscious by boyfriend – intubated by paramedics but remains paralysed

Working out


1. Sinus tachycardia
2. RAD
3. Broad QRS (RBBB morphology)
4. Prolonged QT interval

Dx = Na channel blockade OD


Mx:

1. Give IV NaHCO₃

PREVENTING DIC



DAMAGE CONTROL = STAGED SURGERY



- **Resuscitative surgery** → sufficient surgery to control bleed / contamination (from orifices) and resect obvious ischaemic structures (e.g. small bowel, limb)
- **Stop cosmetic surgery** (e.g. stomas, generating anastomoses) → just stable organs up e.g. packing → minimise abdo compartment syndrome, infection
- **Complete definitive surgery OR later reconstruction** (e.g. suture)

MASSIVE TRANSFUSION PROTOCOL (MTP)

Indications	Target / Goals	Complications of MTP = fever, chills headache										
• pRBC = gold standard = Minimise crystalloid (e.g Hartmann's, NS) • hydrocort 200mg prior to blood transfusion to prevent ADR Group (O neg) to replace: • 50% BV loss in 3hrs • 100% BV loss in 24 hrs <table><tr><td>No trauma</td><td>4u pRBC + 2U FFP</td></tr><tr><td>Trauma</td><td>PLUS TXA</td></tr><tr><td>Fibrinogen < 1g/L</td><td>PLUS Cryoppt (clotting factors)</td></tr><tr><td>Plt < 50</td><td>PLUS Plts</td></tr></table>	No trauma	4u pRBC + 2U FFP	Trauma	PLUS TXA	Fibrinogen < 1g/L	PLUS Cryoppt (clotting factors)	Plt < 50	PLUS Plts	• Temp > 35 • ABG = >7.2, BXS >-6, Lactate < 4 • Ionised Ca = >1.1mM • Plt = > 50 • Hb = depends on haem status • INR ≤ 1.5 • PT/APTT = <1.5x normal • Fibrinogen > 1g/L • Restore perfusion • Correct cell hypoxia • Prevent MOF • Support Haemostasis • Avoid A/E		Acute	Delayed
No trauma	4u pRBC + 2U FFP											
Trauma	PLUS TXA											
Fibrinogen < 1g/L	PLUS Cryoppt (clotting factors)											
Plt < 50	PLUS Plts											
Immune		Haemolytic (DARK URINE)	Haemolytic									
		TRALI → ARDS	Alloimmunisation									
		Febrile non-haemolytic (MOST COMMON)	Posttransfusion purpura									
		Allergic	GVHD									
Non-Immune		Bacterial contaminate (ESP, PLT TRANSFUSION WHICH ARE KEPT AT ROOM TEMP_	BBV (hep B/C/HIV)									
		Circulatory overload (TRACO)	Fe overload									
Don't miss		Electrolyte (hypo/hyperk) (HypoCa = citrate in soln increases excretion of Ca)										
		Coagulopathy										
		Hypothermia										

When to give blood products?

Whole blood (pRBC + FFP) = 530mL (235mL + 295mL)

Product	Dosage (paediatrics)	Indication
pRBC • O neg = universal donor; • AB positive = universal acceptor	1 unit/bag (300mL) = 10g/L rise required. • 235mL of 530mL • Raises K and lactate • Decreases Ca	Hydrocort 200mg prior to blood transfusion to prevent ADR • Hb <100 and high risk of myocardial ischemia • Severe anaemia (e.g. Hb<70) • Major active bleeding and Hb<100
Plts	10mL/kg (1 unit ~ 60mL. 1 pooled bag = 5 units) • 295mL of 530mL • Raises K and lactate • Decreases Ca	• <10 • <20 and high risk (fever, neutropenia, antibiotics, risk of intracranial haemorrhage) • <50 and active bleeding or requires invasive procedure • <80 and requires neurosurgery or ophthalmic surgery • Plt function defects and bleeding (regardless of plt count)
FFP • AB is the universal FFP donor, (no antibodies in plasma)	10 – 20 mL/kg (1 bag ~ 230mL)	• INR >1.5 PLUS needs invasive procedure • INR >1.5 PLUS actively bleeding (e.g. massive transfusion protocol, post-bypass surgery)
Cryoppt	5-10 mL/kg (1 bag ~ 20mL) contains ➢ Fibrinogen (factor 1) ➢ Factor, 8, 13 ➢ VwF and Fibrinectin *Thawed FFP then extracting ppt	• fibrinogen <1.0 and actively bleeding (e.g. massive transfusion protocol) • DIC (main indication) • hereditary hypofibrinogenemia, • haemophilia, • VWF disease
Factor 7	90 mcg/kg	• rescue therapy if ongoing haemorrhage despite correction of pH and temperature, blood products to correct coagulopathy and no clear surgical cause
Prothrombin (F 9) complex:	1mL/kg (25units/kg)	• warfarin overdose (alternative to FFP)
TXA acid:	100mg/kg then 10mg/kg/hr	Massive haemorrhage (PPH, Menorrhagia, GIB)
Granulocyte concentrate		Neutropenic sepsis

D – DISABILITY – COMMON PRESENTATIONS #1

- **Anatomical (FRONT to back)**– sinusitis, eye (glaucoma), scalp tenderness (GCA), brain infection, raised ICP (IIH, SoL), neuralgias, neck-related (referred pain), ear-related (otitis media), muscle related
- **Red-flag** – stroke, bleed, infection (fever), sudden onset headache, LOC
- **Non-related** – migraines, TTH, cluster

	Headache	CNS infection	Seizure
Hx	- Sudden, exertional onset - New headache in > 50 yo - FND (vision, speech, weakness, sensation) Check location and acuity/time-course - Eyes, - slit-lamp - Ears - otoscope	- New onset headache - Fever, rigors, chills - N/V - Photophobia - Neck stiffness (can be just mild) - FND - Confusion / encephalopathy - Non-blanching maculopapular rash - Kernig's and Brudzinski's sign Have they been taking any ABx? Are there any leg pain?	Collateral hx - How were they before? - Prodrome – triggers, systems review - Event – witness, situation, WILD Sx - Post-event – recovery time, symptoms
RF	Pro-coagulant states (e.g. pregnancy, SLE, cancer, sarcoid, vasculitis) Meds (blood thinners, immune suppressors, recent ABx) Illicit drug screen (stimulants e.g. cocaine)	Prodrome illness & ENT infections – URTi, otitis media, orbital cellulitis Immunosuppressed (T2DM, chemo, ATSI) Partially treated infection – persistent headache despite ABx from GP - Trauma - Age - Recent Procedures	- Hx of epilepsy - Recent illness - Known intracranial lesion
DDx	- ICH, SAH - CNS infections - GCA - palpate TMJ - Malignancy Eye pathologies (Optic neuritis, acute angle closure glaucoma), IIH Trigeminal neuralgia - Vertebral artery dissection → palpate vertebral artery sinuses - Cerebral venous thrombosis	Meningitis – B/V/F (MOSTLY viral) children - Strep, pneumo, Neisseria, Hib, GBS Elderly → strep, Neisseria, gram - ve Encephalitis – B/V/F Brain abscess Epidural abscess DDx: - may present asymptotically Meningococcal sepsis (bacteria in blood without any signs of meningism) Meningitis (infection of meninges without non-blanching purpuric rash)	BATH TIME - Brain lesion - Alcohol/drugs withdrawal - Trauma - Hereditary - Toxins - Infection - Metabolic (BSL, Na, K) - Epilepsy DDx of reduced LOC (AEIOU – TIPS) - Acidosis / alcohol - Epilepsy - Infection - Overdose - Uremia - Trauma - Insulin - Psychiatric - Stroke Other DDx: - Syncope (cardiogenic, vasovagal) - Psychogenic - Febrile convulsion (6/12 – 6 years) - Dehydration
Ix	Bloods – FBC, EUC, LFT, CRP, ?INR Imaging - CT within 6 hrs to exclude SAH - MRI - Carotid artery doppler - Slit-lamp examination - LP	Bloods – FBC, EUC, LFT, CRP, cultures Imaging – brain CT/MRI	ECG – monitor source of seizures Urine B-HCG Bloods – FBC, EUC, CMP, VBG (lactate), Prolactin, BSL, B-HCG MRI EEG
Mx	Neurosurgical consult Decompress bleed	ABCDE EARLY Empirical ABx – IV 2g (50mg/kg) ceftriaxone - Bloods - FBC, EUC, LFT, CRP, cultures M/C/S - Dexamethasone within 1hr of ABx - CT before LP (avoid tonsillar herniation due to intracranial mass) Consider - Benzylpenicillin - Acyclovir - Vancomycin - Tacozin	Crowd control → ABCDE - A – maintain patent airway and keep patient on side (jaw thrust, Guedel) - Make sure to expose everything!!! Most fits/seizures self-resolve. If seizure > 5mins then consider medical treatment 1st line = BZD 2nd line = Repeat BZD 3rd line = Keppra or phenytoin 4th line = give what was NOT given already 5th line (infusions) = midazolam, propofol, thiopentone Safety netting - No driving - NO operating heavy machinery - Swimming - Baths - Caution with heights

A Case

JASMINE

- 32 year old – Severe headache
- 1 week post partum
- G5P5
- Obstetric Hx complicated by severe HT
- C sec for HT at 34 weeks
- Epidural for C sec
- D/C for 4 days
- Since last day in hospital has had nausea, feeling generally unwell
- Worsening severe headache

A Case

MOHABIK

- 32 year old male** - Headaches for 1 week
- Moderate, not worst headache ever
- Wakes at night with pain
- Wakes at night to vomit
- Pain worse in the morning
- No neurology
- Pain completely resolves with paracetamol

A Case

AJ

- 8 year old boy
- Fevers 1/7
- 1st Seizure at home self terminated
- Seizure in ED
- No PHx,
- 1 uncle with epilepsy
- No meds
- No allergies
- IUTD

	CVA	ICH	Vertigo/Dizziness
Sx	➤ LOC ➤ FND	Sudden onset FND, reduced LOC, photophobia Headache (thunderclap, exertional)	Is it vertigo or dizziness? ➤
RF	• Vascular-pathology – T2DM, anti-coags	• Trauma • Meds - Anti-coag usage	
DDx	➤ Stroke / TIA ○ Ischaemic ○ Haemorrhagic ➤ Hypo/hyper electrolytes, BSL ➤ Epilepsy / seizures ➤ MS, Migraine ➤ Intracranial infection / mass ➤ Syncope	➤ Stroke /CVA ➤ Syncope	DDx for dizziness ➤ Head, heart, vessels, systemic DDx for vertigo • Peripheral – BPPV, Vestibular Neuronitis, Meneire's, Stroke • Central – Mass, ICH/CVA, MS, Migraine, meds
Ix	• Bloods – FBC, EUC, LFT, CRP Imaging • CT within 6 hrs to exclude SAH • MRI • Carotid artery doppler • Slit-lamp examination	• Vitals • ECG + CXR • Bloods – FBC, EUC, LFT, CRP, COAGs, Group and hold Imaging • CT brain ○ Non-contrast (if after 6 hrs) ○ Contrast (if suspected SAH) ○ CT arch CoW (aneurysm?) ○ CT arch from neck (vertebral dissection) • MRI • Carotid artery doppler	• Head impulse – vestibulo-ocular reflex • Nystagmus ○ Central (absent, vertical, continuous and bidirectional) ○ Peripheral – present, horizontal, unidirectional • Test of skew – cover/uncover test • Dix-hallpike • Imaging – CT vs CT CoW vs MRI • Bloods – troponin, septic screen, VBG, BSL
Mx	Refer to local guidelines based on Mx 1. Call stroke team urgently 2. Urgent Imaging → CT, CTA, perfusion 3. Thrombolysis if within 4.5 hours 4. Clot retrieval (salvageable tissue depends on size of vessel occluded) Non-urgent intervention ➤ MDT ➤ Monitor BP ➤ Aspirin ➤ Anti-coags (e.g. heparin)	SAH ➤ Support – maintain normal EUC, O2, temp, glucose ➤ Maintain BP 90-140 ➤ Reduce ICP – analgesia, HOB 30 deg, anti-emetics for N/V ○ Consider mannitol, hyperventilation if unresponsive ➤ Consult neurosurg ICH ➤ As above with NO strict BP control SDH (elderly fall – check GCS) ➤ Consult neurosurg ➤ More time as usually more stable (do not suddenly pop) EDH (period of lucidity) ➤ Urgent – risk of rapid deterioration ➤ Consult neurosurg for rapid decompression (burr-hole washout)	1st line = Sx control ➤ Prochlorperazine ➤ Promethazine *check for allergies Treat cause ➤ BPPV – Epley ➤ Vestibular neuronitis – prednisone ➤

A Case

21 year old Sebastian presents to country hospital

- Severe sudden onset headache, BP 140/90, PR 115
- Take and Hx and Ex

Manage this patient, DDx

- Patient transferred to neurosurgical team big city center, BP 130/85, PR 95
- Every imaging you can think of CTs, MRI, body scans, contrast, PET
 - Small lesion basal ganglia, possibly slow growing malignancy, ?calcified bleed from birth

Manage this patient, DDx

- Managed on wards for headaches, BP 160/95, PR 95
- Agitation, request hot showers, family aggressive and confrontational
- Lignocaine infusion for headache, mental health team involved
- TF back to country hospital

Manage this patient, DDx

- Unable to get repeat imaging TF to slightly less country hospital with the ability to MRI
- Sweaty, BP 170/100, PR 120, afebrile, request hot shower

Manage this patient

D – ULTRASOUND TIPS #1

SCENARIO	QUESTIONS	USS ACTIONS
CARDIAC ARREST	- Is it reversible? (4 H's and 4 T's) - Are both lungs ventilating	- Heart (tamponade, LVH, or reduced contractility) - Lungs - Elsewhere - IVC, AAA, free fluid
SHOCKED PATIENT	- Why is patient in shock? - Do they need fluids?	- Anterior lungs – wet, dry, chunky? - IVC – full or empty - Heart - Elsewhere - IVC, AAA, free fluid
BREATHLESS PATIENT	Why are they breathless?	- All of the lungs – wet, dry, chunky? - IVC - Heart - Scan for DVT – upper femoral/lower femoral, popliteal or below knee

C	Continue Compressions and Echo
O	Oxygen Away
A	All Else Stand Clear
C	Charge Defibrillator
H	Hands Off Compressions
R	Rhythm Check/Record Echo
E	Echo Off
D	Deliver Shock <i>OR</i> Dump Charge

7 GOLDEN RULES FOR USS

- 'Resus-only'** (critically unwell: shocked / breathless / peri-arrest to elicit US signs of some diseases e.g. massive PE or pneumonia)
- Clinical Diagnosis BEFORE US** (e.g. FBC with Hb = 4)
- Only ask questions that you can answer**
- Repeat scans are crucial (change over time)**
- USS are NOT 100% accurate
- When in doubt, be a doctor (clinical picture > USS findings)**
- A fool with a stethoscope will be a fool with an ultrasound**




WHICH PROBE TO USE?

			
Probe type	Curved (curvilinear) array	Linear array	Sector (Phased) array
Indication & preset	Deep tissue visualisation Lung (pneumothorax, effusion) Heart (subcostal window) Abdomen (abdo, eFAST, AAA, early pregnancy) IVC	Superficial tissue visualisation - Lung pleural surface ONLY Vascular access (vascular / venous preset)	Lung (pneumothorax, effusion) Heart (subcostal window) Between ribs (Cardiac present – Nb: image wrong way around)

TIPS TO IMPROVE IMAGE







- Use **MORE gel** (reduce reflection at surface of skin)
- DARK** room
- Correct patient position
- Right probe and preset
- Correct probe orientation
- Set depth (deeper than structure you're scanning)
- Set adequate gain (fluid just appears black – to differentiate between fat and fluid)
- Highest frequency = better resolution (low frequency = for obese patients)**

Interpretation of images:

			
Bite in GB = black (anechoic) Below brighter = posterior acoustic enhancement	Stones in GB (HYPER echoic) Casts posterior acoustic shadow	Ascites (simple fluid) (kidney is below)	SC abscess
Curved probe	Curved probe	Curved probe = Paracentesis – watch out for - Inferior epigastric arteries - diaphragm	Linear probe

BEWARE OF anisotropy (esp. nerves) → specific orientations to better visualise structures

D – ULTRASOUND TIPS #2

		
Lung USS	Cardiac USS	Vascular Access
Curvilinear	Sector probe (to fit b/w ribs) ➤ subcostal position	➤ Linear probe ➤ Use BP cuff as tourniquet ➤ Always use LONGER cannulas
• Pneumothorax (NO lung sliding) • Fluid/effusion (image 2) ○ B lines ○ Rockets and comets	For critically unwell patients 1. Check Heartbeat 2. Effusion → ?tamponade 3. RV bigger than LV → big PE 4. Normal → may need move volume	➤ Cannulation of difficult veins
		 Q8. This subcostal image (taken during CPR) shows: A. cardiac standstill CORRECT B. tamponade FALSE C. RV compressing LV FALSE D. Other FALSE
	 Q9. Arrested patient. This apical image shows: A. cardiac standstill FALSE B. tamponade CORRECT C. RV compressing LV FALSE D. Other FALSE Tamponade	Cardiac Standstill

In ALS → Use either curvilinear or sector probe (whichever convenient) → aim to rule out 4 H's and 4 T's

D – MANAGING DIFFICULT SITUATIONS → “SPIKES”

Setting

- Turn off phone
- Sit on chair
- Make sure the patient is with a supportive person
- Avoid presumptions

Consider your patient

- Know their name!
- Review their notes prior to any discussions, know as much as you can know


Consider your audience

- Are you talking to a patient?
- Know who you will be addressing if possible before meeting

Consider your news

- Is it a complete shock?
- Should you use a “warning shot”. Tell them what you want them to know first, before anything shocking
- People will forget everything you said after the bad news breaks → **need to see the word death or your mum/dad/son is dying**

SPIKES PROTOCOL



: On the phone

- Do not say patient had died on the phone
- Please come in as ____ is critically unwell.

Finishing up

- Offer social worker support
- Give time to process

E - TRAUMA: Primary Survey

What is a trauma patient?


- **High impact** = MVA, MBA, Plane crash,
- **Penetrative** = Stabbing, shooting
- Significant fall (esp. elderly)

What do I consider?


- Collateral hx (ambos, police, bystanders)
- AMPLE
 - Allergies, meds, PMHx (diabetes, vascular, collagen diseases),
- What happened before event? During trauma? And after? (i.e. what was given as Rx)

Groups	Special Considerations	Intervention
Children 	- Leading cause of death - Immature, anatomic/ mechanical features - Vigorous physiologic response - Limited physiologic reserve - Consider: <i>Size, dosage, equipment, SA, and psychology</i>	A: Larynx anterior and cephalad, short tracheal length B: Chest wall pliability (contusions may not present) C: Vascular access, fluid volume, vital signs, and urinary output D: Vomiting, seizures, and diffuse brain injury MSK: Immature skeleton, fracture patterns
Pregnancy 	- Anatomic/ physiologic changes modify response to injury - Need for fetal assessment 1st Priority: Maternal resuscitation ○ Identify early shock ○ X-ray mother	- Gestation and position of uterus - Physiologic anaemia ↓ Pco2 ↓ Gastric emptying - Supine hypotension - Isoimmunization (Anti-D for Rh -ve mothers) - Sensitivity of fetus
Elders	5th leading cause of death - Diminished physiologic reserve and response <u>Co-morbidities:</u> Diseases/ Meds	
med	- BB may hide tachycardia Intoxicated	Drug screen Safety of myself and others
athletes	normal HR when actually tachycardic	Low threshold to admit to ICU

PRIMARY SURVEY



MULTI TRAUMA ACUTE MANAGEMENT




BEWARE: Trauma-induced coagulopathy (TIC)

- **Acute traumatic coagulopathy (ATC)**
 - Hyperfibrinolysis
 - Fibrinogen loss
 - Plt dysfunction/disaggregation
- **Resuscitative coagulopathy (iatrogenic)**
 - Due giving too much fluid free of coagulation factors
- **Prothrombotic phase**

Solution

- **Reduce crystalloids** → 1:1:1 ratio → RBC:plt:plasma
- **Permission hypotension** (i.e. accept low borderline BP)
- **Damage control surgery**
- **Blood & coags**



E - Secondary Survey – “Top to toe”

History	Exam	Do the following:
A Allergies M Medications P Past illnesses / Pregnancy L Last meal E Events / Environment GCS Scalp Ears (incl TMs) Eyes (incl pupils/acuity/fundi) Facial bones Mouth Neck C-spine Clavicles Chest wall Chest movement Breath sounds Heart sounds Abdomen Pelvis Hips Thighs Knees Legs Ankles Feet Shoulders Upper arms Elbows Forearms Wrists Hands Fingers Back (incl spine) and flanks (log roll) Perineum Genitalia Rectal exam Urinalysis	- Head → ENT → Neck (uncover any c-cover) - Chest - - Abdo - Pelvis - Extremities (upper → lower) Seek the following: Bleeding Swelling (incl. haematoma) Tenderness Lacerations (incl. entry/exit) Deformity (bones) Discolouration Crepitus (incl. subcutaneous) Pulse - Ischaemia (limbs) Functional impairment - Neurological (power, tone, reflexes) - Viscera (lungs, heart, GIT) - MSK	1) Stop bleeds - IVF, - pressure on wound, - pelvic binders, - reduce fractures with traction 2) Stop pain = Analgesia (IV morphine) 3) Stop infection - ABx or tetanus prophylaxis, - Sterile pads on wounds Adjunctive things to do: - Photo major wounds - Bedside → UA, ECHO - Bloods - Imaging → X-ray, CT - Invasive → Bronchoscopy, oesophagoscopy Do NOT delay transfer to 3° hospital (trauma centre) – bring relevant imaging and results

***NB: Tertiary survey avoids MISSED injuries → (repeat 1°, 2°, relook at imaging/tests etc.)**

- Within 24 hrs** (repeat primary and secondary survey and any imaging) → PROCEED TO NEW IMAGING
- DO NOT MISS** → Mild TBI, PNI, C-spine #, bowel injury, knee ligament ruptures, acromioclavicular dislocation

Trauma Triaging

Multi-tasking: Do it once and do it correctly!


- Get line in and bloods prior to hx – get ALL the tubes
- Prepare lubricants for DRE for PR bleeding
- Vag speculum exam for PV bleeding

Causes	Good trauma system	Trimodal Death distribution
Blunt (road accidents, fall, blast) Penetrating (stab/impale, gunshot) Burns Electrocution Chemical Combination of above - Don't be distracted from the occult → what is most life-threatening?	- Minimise time from injury to care - Prioritise patient's threat to life and disability Preparation crucial - Get team ready & wear protective equipment (e.g protect against BBV – Hep B/C, HIV) Timeframe for multi-trauma: 1st hr = critical initial assessment & resus 2nd hr = definitive care Treat the cells! → STOP ISCHAEMIA (hypoperfusion) → INFARCT (cellular response to ↓perfusion)	

TRAUMA SERVICES PLANNING	REGIONAL TRAUMA CARE SYSTEMS	TRAUMA SERVICE COMPONENTS	Trauma Team
- Population - Trauma frequency - Existing resources - Transport times - Geo-political issues <u>Major Trauma Services/Hospitals</u> (e.g. John Hunter, St George, RPA)	Pre-hospital services - Transport - Communication Acute hospital services Rehabilitation services Interhospital transfer processes Specialist centres: - Burns - SCI	Need good communication b/w: - Emergency Medicine - Surgery - Intensive Care/Anaesthetics - Operating Suite - Radiology - Blood Bank - Allied Health - Rehabilitation	- Emergency Medicine Registrar 3 RN's from Emergency Department - ICU Registrar - Trauma Fellow - Trauma Nurse Co-ordinator - Surgical Registrar - Radiographer - Social Worker - Blood bank on standby

What do I do?	PREHOSPITAL TRIAGE (MIST)	Activation Criteria for T/F to MAJOR TRAUMA Centre: "HANDOVER"
PREHOSPITAL TRIAGE – who goes where? Hospital? Who needs the Trauma Team? Mass casualty? – who gets the resources? 1st half of anybody's life – more likely to die from injury (i.e. road crash, burns, drowning) than from any other cause > 65 yo = Most deaths due to falls and suicide	- Patient's name, age and sex Mechanism - All Penetrating - MVA - fall from standing height Injuries (obvious) - Beware low impact injuries for elderly Signs (Vital) - Pulse, BP, RR, GCS (at scene, on arrival and current) Treatment required - IVF, ABx (tetanus?), Analgesia - FBC, EUC, COAG, x-match - XR, CT, ECG, ECHO, FAST	(1) Introduce self and location (2) Brief History of patient and Injury: - MVA ejected from vehicle/fatality - Pedal cyclist, motor cyclist or pedestrian hit by a car or truck - Fall from height > 5 m - Inter-hospital transfer (3) What were the injuries?: Open/closed Fracture to ≥2long bones Soft tissue injury Bleeding Crush injury or limb amputation Any Penetrating injury (e.g. PT, haemothorax) Burns to airway or smoke inhalation (or > 15% BSA adults, > 10% BSA children) (4) Vitals and What was done? - A = adjuncts - B = HM, Bag-valve, Ventilator - C = Fluids, peripheral stimulation - D = GCS, c-spine - E = splinter? Reductions?

Trauma Day – Fractures and Splinting

Hx	Exam	Ix	Describing fractures
AMPLE Co-morbidities (DM, collagen and vascular disease) Events (before, during, after) <i>E.g. broken collarbone + hidden c-spine injury</i> 	Give analgesia 1st - Vitals Look at EVERYTHING - Site <i>e.g. femur #</i> → <i>assume 1L blood loss</i> - Deformity - check for OTHER wounds (e.g. punctures) - above and below joint Feel - vascular – distal pulses - neuro – sensation Move - ROM – passive vs active - Pain out of proportion (compartment syndrome) Measure (mark out) - Swelling - Limb length - Special tests	Bloods - FBC (Hb) - VBG (lactate and Hb) Group and hold (if sig. bleed) - EUC, LFT – baseline if OT required Imaging - XR (may miss fractures!!) 1. Check cortex. 2. Bony alignment/dislocation 3. Lipo-haemathrosis 4. Avulsion # - fragments of bone pulled off by tendon/ligaments 5. Fat pad (sail sign – esp. elbow – blood pushing away bone) 6. Effusions 7. Salter-Harris # in kids 8. Check above and below - USS (? If foreign body, fluid) – use lots of gel to minimise need to apply XS pressure - CT – if not sure on XR (esp. wrist, comminuted #, scapula, ribs) - MRI (ligaments, scaphoid #, knee)	Pt name, gender, age, incident, dominant hand and occupation (<i>e.g. pianist, surgeon</i>) Open/closed Location (left/right vs proximal/distal/mid) Fracture type (oblique, TV, spiral, greenstick, impacted, comminuted) Angulation (direction of tilt of distal fragment) Fluid? Neurovascular intact? AMPLE – fasting status Health insurance General Mx - Plaster (e.g. Sydney sandwich & backslabs) - Avoid dirty objects with open wounds - Avoids objects that splinter


			
Anterior Fat pad sign of supracondylar fracture	Closed Spiral L) mid-humerus #	Closed Comminuted L) proximal-mid femur with medial displacement	Wrist injury with Dorsal displacement of wrist bones (empty cup = lunate bone = "crescent moon sign")

Trauma Day – Trauma Wounds & Minor Injuries

Hx	Exam	Ix	General Mx
AMPLE Co-morbidities (DM, collagen & vascular disease) Events (before, during, after) (fall, bite, penetrating object) Is wound dirty or clean? (is it from sterile place – water, dirt, raw chicken, FB, cats/dog bites) Associated injuries? (how much bleeding?, sensation loss, SOB cosmetic issues for young women – nipple) Occupation – impact on work (e.g. surgeon, driver) 	Give analgesia 1st - Vitals Look - Site & depth - shape & borders - exudative (infective) - erythema (oedema) - colour Feel - tenderness (bone, soft tissue) - warm to touch - neuro – sensation - vascular – distal pulses Move - ROM – passive vs active - Joint above and below Pain out of proportion (beware sown up wound) Measure (mark out) - Depth and size	Vitals Bloods - FBC (Hb) - VBG (lactate and Hb) - CRP - EUC, LFT and CMP (if ABx req) ?Septic screen (blood cultures) Imaging - XR - EXCLUDE fractures - Osteomyelitis - Foreign body - USS (? If foreign body if NOT radioopaque) - CT (details of bone) - MRI (ligaments) Reference: - Plastics (e.g. nipple wound) - Ortho – hand surgeon	- Good EARLY cleaning w/ 0.9% NS → <i>remove any MSSA, MRSA</i> If very painful → use local LA in 50mL syringe (avoid injecting into arterioles) Remove any clot/dirt to see the wound w/ cleaning Stop any bleeding - Clean to find source of bleed - apply pressure with gauze - Adrenaline - Haemocoagulant dressing - AgNO₃ sticks or cautery Keep wounds covered with suitable dressings Types of dressings (definitive care) Wet (w/ 0.9% NS) – non-stick, duoderm (ulcers), mepilex, bactigras (paraffin based – create nice seal) Dry – Melolin Padding – Combine (Secure with crepe (E.g. turban for scalp bleed)

Trauma Day – Trauma and Burns

Hx	Exam + general Mx	Additional Mx								
• AMPLE • Co-morbidities (DM, collagen and vascular disease) • Events (before, during, after) ◦ ISOLATED trauma (house fire) ◦ Polytrauma burn + other injuries (car accident/car fire or explosion) ◦ Substance inhalation (cyanide due to burning plastics, CO - soot) • Duration of burn • Mechanism ◦ Domestic steam (<100°) ◦ Industrial steam (>100°) ◦ Fat (>250°) ◦ Open flame (>500°) ◦ Electrical 240V ◦ Chemical)	1. Personal and staff PPE + decontaminate patient 2. Remove items (clothing, jewellery) as they retain heat 3. Cool burn (cool running water → 20 mins) 4. Simultaneous ABCD <table><tr><td>A</td><td>Check for signs of airway burn → oedema → req early intubation (<i>they get worse before getting better</i>) ➢ Singed facial hair (loss of eyebrow, beard, nasal hair) ➢ Soot in airway ➢ Beware of sore throat or Hoarse voice </td></tr><tr><td>B</td><td>Inhalation injury may extend into trachea and lungs ➢ Causes local/systemic inflammation → ARDS ➢ High mortality rate → consider Asthma, COPD, existing lung disease ➢ Rx: respiratory support </td></tr><tr><td>C</td><td>Burn affects circulation in 2 ways 1. Systemic inflammation = oedema and loss of intravascular volume (hypovolemia) → Aggressive IVF – Parkland Formula a. Calculate %TBSA → rule of 9's 2. Burn depth a. Burn erythema – bad sunburn b. Partial thickness c. Full-thickness d. Circumferential Bloods → Check ABG → for CO poisoning </td></tr><tr><td>D+E</td><td> • GCS, BSL, Analgesia (<u>GCS affected by ketamine, sedation</u>) ◦ Check movement / sensation (? SCI) • Thermoregulation → use warm fluids, dressing, and warm with hair hugger 40°C (burn patients quickly become cold) • Moisture Regulation → dressing (reduce evaporation to minimise hypovolaemia) </td></tr></table>	A	Check for signs of airway burn → oedema → req early intubation (<i>they get worse before getting better</i>) ➢ Singed facial hair (loss of eyebrow, beard, nasal hair) ➢ Soot in airway ➢ Beware of sore throat or Hoarse voice	B	Inhalation injury may extend into trachea and lungs ➢ Causes local/systemic inflammation → ARDS ➢ High mortality rate → consider Asthma, COPD, existing lung disease ➢ Rx: respiratory support	C	Burn affects circulation in 2 ways 1. Systemic inflammation = oedema and loss of intravascular volume (hypovolemia) → Aggressive IVF – Parkland Formula a. Calculate %TBSA → rule of 9's 2. Burn depth a. Burn erythema – bad sunburn b. Partial thickness c. Full-thickness d. Circumferential Bloods → Check ABG → for CO poisoning	D+E	• GCS, BSL, Analgesia (<u>GCS affected by ketamine, sedation</u>) ◦ Check movement / sensation (? SCI) • Thermoregulation → use warm fluids, dressing, and warm with hair hugger 40°C (burn patients quickly become cold) • Moisture Regulation → dressing (reduce evaporation to minimise hypovolaemia)	Take photo of burn (with consent) • 1st aid Dressing (e.g. glad plastic wrap – single sheet) 1) Deroof blisters 2) Cover all burn 3) Barrier to infection 4) Improve pain and temp. regulation as prevents air touching burned skin • Analgesia (IV morphine) • ADT (tetanus) • IDC -measure UO • +/- NGT - if intubated • Avoid ABx – causes harm in burns Transfer to definitive care – refer EARLY • Dedicated burn units = Concord (non-trauma) RNSH (adult + trauma), Westmead (paeds, adults + trauma) • Inpatient 1) Adult > 20%, children > 10% 2) Facial / intubated 3) Electrical/ chemical 4) Suspected NAI • Outpatient 1) Full thickness of hand, feet, perineum 2) Does NOT need inpatient Mx • Depth of burn can last for 7-14 days depending on initial Rx (first aid + resus)
A	Check for signs of airway burn → oedema → req early intubation (<i>they get worse before getting better</i>) ➢ Singed facial hair (loss of eyebrow, beard, nasal hair) ➢ Soot in airway ➢ Beware of sore throat or Hoarse voice									
B	Inhalation injury may extend into trachea and lungs ➢ Causes local/systemic inflammation → ARDS ➢ High mortality rate → consider Asthma, COPD, existing lung disease ➢ Rx: respiratory support									
C	Burn affects circulation in 2 ways 1. Systemic inflammation = oedema and loss of intravascular volume (hypovolemia) → Aggressive IVF – Parkland Formula a. Calculate %TBSA → rule of 9's 2. Burn depth a. Burn erythema – bad sunburn b. Partial thickness c. Full-thickness d. Circumferential Bloods → Check ABG → for CO poisoning									
D+E	• GCS, BSL, Analgesia (<u>GCS affected by ketamine, sedation</u>) ◦ Check movement / sensation (? SCI) • Thermoregulation → use warm fluids, dressing, and warm with hair hugger 40°C (burn patients quickly become cold) • Moisture Regulation → dressing (reduce evaporation to minimise hypovolaemia)									

			
Burn erythema (1 st deg)	Partial Thickness (2 nd deg)	Full thickness (3 rd deg)	Circumferential Full thickness
Bad sunburn Painful - No scarring Rx: aloe vera!, burnaid	- Blisters (less deep) - Cherry red (more deep) Blanches Hair Intact Painful (dermal pain nerves injured)	- Pale - Feels like leather, does NOT blanch, hair loss Painless (cutaneous nerve "killed")	ALL the way around - Loss of dermal compliance (flexibility) Skins becomes like a tourniquet causing ischaemic limb, SOB (torso)

Trauma Day – BURNS calculating resuscitation fluids


Steps

- Calculate %TBSA** – Adults ≥ 20% or child ≥ 10%
- Parkland's formula** (see below)
- IVF – Hartmann's** (to minimise K loss)
- Check IDC – urine output**

2-4ml x (ideal) weight (kg) x %TBSA
 - 50% in first 8 hrs
 - 50% in next 16 hrs
 titrate to urine output – adult = 0.5ml/kg/hr

What happens if it does not work? Eschar?

- Eschar** = tough, inelastic mass of burnt tissue (secondary from full thickness circumferential burns)
- Compartment syndrome** = eschar + swelling (i.e. circumferential burns)
 - Vascular compromise of limb (distal ischaemia)
 - Impending / actual respiratory compromise
- Treat with escharotomy**
 - Life-saving procedure – improve blood flow
 - Incise tissue along predefined escharotomy lines



Rule of 9's:

***ONLY** includes 2nd and 3rd degree burns

**Not used for children (which uses rule of 5's)

Case Study: Acute Trauma:

1. A 26 yr old healthy man was riding a motorcycle which collided with an SUV. He has just been brought to hospital by ambulance. He seems alert, appears pale and distressed and is complaining of severe pain in his R leg. His vital signs are P 120, BP 100/80, RR 20/min. The ambulance officers report that he has swelling and deformity of his right thigh and that his right foot is pale and cold. He also has laceration to his L upper arm, to which a dressing and bandage have been applied to stop bleeding.

DDx	Mx
- Haemorrhagic shock secondary to Shaft of femur # - Nb: NOF# = Shortened leg + ER	Get help – surgical reg A – patent B – breath sounds, FiO2 (NP) C – IV Access (2x large bore cannulas) - IVF – 500mL 0.9% NS - Bloods – FBC, EUC, LFT, COAG + GROUP + X-MATCH - IV Analgesia – morphine - IV anti-emetics –
Key findings	
- Hypotension – hypovol. - Tachycardia – compensating for hypoTN to maintain CO R) Pale/cold foot → ischemic foot (risk of amputation) - Vascular injury = Suggests bleed from superficial femoral artery which drains down into popliteal artery	D – GCS and BSL E – Secondary survey → hypothermia (need warm blankets) ➤ Traction on leg to straighten shortened segments of femur (prevents ischaemia and compartment syndrome) ➤ Angiography + stent OR ORIF (restore femur alignment) ➤ Maintain pressure on bandage dressing until haem stable

2. A 18yr old young woman was crossing the road when hit by a car travelling at about 50km/hr. She was thrown sideways and impacted the guard rail on the side of the road. She has been complaining of abdominal pain during her ambulance transport to the hospital. En route she has been a bit confused, BP has fallen from 110 syst to 90. She looks quite pale and her hands are cool and clammy while her vital signs are P130, BP 85/60, RR 28/min. Her airway is clear and her chest looks and sounds normal (i.e. equal breath sounds on both sides) but her abdomen appears a bit distended and is quite tender across the upper region.

DDx	Mx	Other Mx
Haemorrhagic shock secondary to: - Haemothorax - Intra-abdo bleed - Long-bone # - Pelvic # - External losses	Get help – resus bay A – patent B – breath sounds, FiO2 (NP) C – IV Access (2x large bore cannulas) - IVF – 500mL 0.9% NS - Bloods – FBC, EUC, LFT, COAG + GROUP + X-MATCH - O -ve (universal donor blood) → while waiting for donor blood D – GCS assessment and BSL E – Secondary survey → hypothermia (need warm blankets) ➤ CXR – tension pneumothorax + multiple rib #, intrapleural bleed ➤ Pelvic XR – hip # and long bone # - Pelvic ring binder ➤ eFAST → assess internal abdominal trauma + pericardial and pleural cavity - mobile mesenteries (transverse, sigmoid, small bowel mesentery) - spleen or liver rupture ➤ Seek additional ICU help → insertion of chest drain	Consider: Urethral bleed → DRE → check prostate position (high-riding) → urethrogram BEFORE IDC or SPC CT angiogram via descending aorta → identify extravasation MTP protocol (pRBC : plt : FFP) → >4 U blood required
Key findings		
Confused ?brain injury ?Cerebral hypoxia ?EtOH/drugs - HypoTN → pale/cool → Vasoconstriction and peripheral shutdown - Tachypnoea - Abdo distension		

3. While removing a fallen tree branch from his roof, a 73 yr old man fell about 3 metres from a ladder and landed on the ground on his right side. He was able to get up but noted sharp pain in his right chest when he breathes. As he began to feel short of breath, an ambulance was called to bring him to hospital. He is alert but anxious and slightly cyanotic around his mouth. His airway is clear but his RR is 30/min, while his other vital signs are P120/min and irregular and BP 100/70. There is a crackling sensation to palpation and auscultation over his right chest and breath sounds are difficult to hear on that side but seem normal on the left side.

DDx	Mx
- Tension Pneumothorax (spot diagnosis) - Rib # → subcutaneous emphysema - Pulmonary Embolism - TBI – EDH/SDH	Get help – resus bay A – patent B – absent breath sounds, FiO2 (NP) C – IV Access (2x large bore cannulas) - IVF – 500mL 0.9% NS - Bloods – FBC, EUC, LFT, COAG + GROUP + X-MATCH D – GCS assessment and BSL E – Secondary survey → hypothermia (need warm blankets) ➤ CXR 1st – tension pneumothorax + multiple rib # ➤ CXR 1st before inserting chest drain (safely) = large 32Fr gauge → drains both and liquid/blood → connect it to underwater sealed drainage (3 bottle system) ➤ Pelvic XR – hip # and long bone # ➤ Non-contrast CT head ➤ Seek additional ICU help → insertion of chest drain
Key findings	
Irregular - ?AF history ?anti-coag = worsens bleed ?BB = hides tachycardia Tension pneumothorax : - Distended neck veins - Deviated trachea Cyanosis - reduced O2 supply - Hypotension – hypovol. Crackle sensation = subcutaneous emphysema (RF for pneumothorax) → NO management needed	

4. Woman aged 33yrs was struck in the head and knocked unconscious by a baseball during a regional competition. She appeared to recover quickly but 10 minutes later she collapsed and has been rushed to hospital. She has a patent airway and is breathing normally. Vital signs are P 90, BP 110/70, RR 15/min. She responds to painful stimuli but not to voice. She has mild swelling over the left temporal area of her head. After Primary and Secondary Surveys, there do not appear to be any other injuries.

DDx	Mx	Other Mx
Extra-Dural haemorrhage (classic presentation of lucid interval following injury) – middle meningeal artery	Emergency !! (as primary and secondary survey completed) ➤ contact Radiology to request urgent non-contrast CT brain ➤ contact Neurosurgical reg and OT staff– update need for OT ready after non-contrast CT brain	What can be done now? ➤ NBM ➤ FiO2 + IV fluids → aim to avoid any hypoxia and brain ischaemia
Key findings	➤ If regional → call tertiary hospital immediately → may receive guidance for burr-hole	➤ Check for dilated pupils / headaches / fluid chart ➤ Check meds for anti-coags/anti-plts ➤ Plan for craniotomy → evacuate haematoma
Direct witnessed TBI + Lucid interval		

